



Maternal Overweight, Obesity and Excess Gestational Weight Gain:

Identification of Maternal and Perinatal Implications
and Primary Maternity Care Providers' Opportunities
for Interventions to Improve Health Outcomes

BC Perinatal Health Program

February 27, 2009

I. Acknowledgements

The research, analysis and writing for this report were prepared by:

Ng, Jennifer, MBBS

Magee, Laura A. ¹⁻⁵, MD, FRCPC, MSc, FACP

Yarker-Edgar, Kristen V. ^{5,6}, MSc, RD

Departments of ¹Medicine, ²Obstetrics and Gynaecology, and ³Population and Public Health, University of British Columbia, Vancouver, BC

⁴BC Women's Hospital and Health Centre, Vancouver, BC

⁵BC Perinatal Health Program, Provincial Health Services Authority, Vancouver, BC

⁶Ministry of Healthy Living and Sport, Province of British Columbia

Supported in part by funding from the Ministry of Healthy Living and Sport, Province of British Columbia

II. Table of Contents

I.	Acknowledgements	2
II.	Table of Contents	3
III.	Executive Summary	5
IV.	Introduction	7
V.	Overweight and Obesity Related to Childbearing	7
	DEFINITIONS	7
	RISKS OF OVERWEIGHT AND OBESITY RELATED TO CHILDBEARING	8
	Fertility	8
	During Pregnancy	8
	Maternal	8
	Gestational Diabetes (GDM)	8
	Hypertensive Disorders of Pregnancy (HDP)	9
	Thromboembolism	9
	Labour	10
	Caesarean Section	10
	Anaesthetic Complications	11
	Postpartum Haemorrhage	11
	Breastfeeding	11
	Fetal and Perinatal Issues	11
	Miscarriage	11
	Perinatal Mortality	11
	Congenital Malformations	12
	Pre-term Delivery	13
	Neonatal Morbidity	13
	Infant Birthweight	13
	The Risks in Perspective	14
	MEDICAL ASSESSMENT CONSIDERATIONS/ CHALLENGES	14
	LONG-TERM MATERNAL AND PAEDIATRIC IMPLICATIONS	14
	COST IMPLICATIONS	15
VI.	Gestational Weight Gain	15
	GUIDELINES	15
	RISKS OF EXCESSIVE GESTATIONAL WEIGHT GAIN	16
VII.	Supporting Women and Health Care Providers to Reduce Risks	17
	INCORPORATE BEHAVIOUR CHANGE THEORY & PRACTICES	17
	CHECKLIST	18
	PRE-PREGNANCY	18
	Healthy Eating, Physical Activity, Substance Reduction and Avoidance	18
	Weight Loss	18
	Folic Acid Supplementation	19
	Optimize Treatment of Co-morbidities	20
	DURING PREGNANCY	20
	Multidisciplinary Support	20
	Screen for GDM	20
	Set an Appropriate Weight Gain Goal	20

Maternal Weight

<u>Encourage Healthy Eating and Physical Activity (to achieve weight gain goal)</u>	21
<u>Healthy Eating</u>	21
<u>Physical Activity</u>	21
<u>Barriers to Healthy Eating and Physical Activity</u>	21
<u>Related to Health Care Providers</u>	21
<u>Related to Women</u>	22
<u>Effectiveness of Lifestyle Interventions</u>	23
<u>POSTPARTUM</u>	24
<u>Prevention of Weight Retention</u>	24
<u>Weight Loss Strategies</u>	25
<u>Outside Pregnancy</u>	25
<u>Tailoring Weight Loss Programs to Women Who are Postpartum</u>	26
<u>Motivations and Barriers to Change</u>	27
<u>Screening for Cardiovascular Risk Factors</u>	27
<u>VIII. Role Modelling to Prevent Childhood Obesity</u>	27
<u>IX. Conclusions</u>	28
<u>Appendix A: Gestational Weight Gain Recommendations</u>	30
<u>Appendix B: Pregnancy Care Checklist</u>	31
<u>Appendix C: References</u>	33

Maternal Overweight, Obesity and

Excess Gestational Weight Gain:

Identification of Maternal and Perinatal Implications and Primary Maternity Care Providers' Opportunities for Interventions to Improve Health Outcomes

III. Executive Summary

The obesity epidemic has become a major public health concern in both industrialized and developing countries. Women of reproductive age are also experiencing the obesity epidemic. In 2005/6, thirty percent of British Columbian women had a pre-pregnancy body mass index (BMI) within overweight (19.9%) or obese (11.4%) categories.

This report reviews: i) the association of maternal overweight and obesity with both adverse pregnancy outcomes and long-term adverse outcomes for mothers and babies; ii) the association of excess gestational weight with both adverse pregnancy outcomes and long-term adverse outcomes for mothers and babies; iii) approaches that can be taken in primary obstetrical care to promote healthy weight among women who are either planning pregnancy, pregnant, or postpartum.

There is a dose-response relationship between maternal weight and adverse pregnancy outcomes, such that risk is: increased with excessive gestational weight gain; greater with pre-pregnancy overweight; and greatest with pre-pregnancy obesity. Women who are overweight or obese are at increased risk for a range of pregnancy complications, including gestational diabetes mellitus, pre-eclampsia, Caesarean section, maternal mortality, and labour induction. Women who are overweight or obese also experience greater risk of infertility compared to women of a lower weight. Women who are obese are at higher risk of complications than women who are overweight, but women who are overweight are still at higher risk than women of normal pre-pregnancy weight. The pregnancy-related risks of obesity are comparable in magnitude to other well-recognized antenatal risk factors, such as smoking or alcohol.

Risks such as stillbirth, fetal death, macrosomia and large for gestational age, neural tube defects, meconium aspiration, and shoulder dystocia are associated with overweight and obesity during pregnancy. Maternal obesity is also a risk factor for childhood obesity and obesity in early adult life.

Pregnancy weight gain within the U.S. Institute of Medicine guidelines (on which the Health Canada guidelines are based) is associated with the best pregnancy outcome for mother and child. However, weight gain alone is not a good predictor of poor pregnancy outcome since the amount of gestational weight gained by women with good pregnancy outcomes is widely variable. Women who are overweight or obese are more likely to exceed IOM guidelines. Risk factors for excessive gestational weight gain include: younger age, being single, ethnicity, current eating habits, self-report of low numbers of

Maternal Weight

supportive individuals in women's lives, and receiving incorrect advice (or no advice) from health care provider(s).

Postpartum weight retention is most strongly associated with excessive weight gain in pregnancy, particularly excessive first trimester weight gain. There is mixed evidence regarding whether postpartum weight retention is associated with pre-pregnancy overweight or obesity. Other risk factors for postpartum weight retention include: unhealthy eating habits, low physical activity, low socioeconomic status/education, low affect/ depression, ethnicity, and self-report of low numbers of supportive individuals in women's lives.

While there is evidence that moderate weight loss will improve fertility, it is assumed that weight reduction before pregnancy will improve future pregnancy outcomes. Outside of pregnancy, recommendations for the achievement of a healthy weight involve lifestyle interventions which include healthy eating, physical activity, and behavioural therapy. However, best practices have yet to be determined as few interventions have succeeded in promoting sustained weight loss, and drop-out rates are high. Lifestyle interventions during pregnancy are aimed at curbing gestational weight gain, not weight loss. Interventions during pregnancy have had mixed results and better practices have yet to be determined. The mixed results of existing intervention studies promoting healthy weights for women pre-pregnancy, during pregnancy and postpartum, emphasize that there are no simple solutions. These findings should serve to motivate the development of better women-centred interventions based on published successes and failures and based on women's motivations and barriers to adopting lifestyles that promote a healthy weight. Because of barriers present when planning a pregnancy and the recognized non-pregnancy related health benefits of achieving a healthy weight, steps should be taken to achieve this healthy weight outside of pregnancy (by promoting weight gain within the IOM guidelines). There should also be postpartum follow-up.

Herein is a checklist of better/ promising practices for primary obstetrical healthcare practitioners to support women who are overweight or obese pre-pregnancy, during pregnancy and postpartum, with a goal to improve maternal and perinatal outcomes.

IV. Introduction

The obesity epidemic has become a major public health concern in both industrialized and developing countries^{1,2,3,4,5,6,7,8,9,10}. In the general population in Canada, the prevalence of overweight has stayed relatively constant while the rate of obesity has risen substantially. In 1970-1971 overweight and obesity rates were 40.0% and 9.7% (respectively)¹¹. In 2004 the rates were 36.1% and 23.1% (respectively)¹².

Women of reproductive age are also experiencing the obesity epidemic. Data from antenatal records in 2005/6 reveal that 30% of British Columbian women had a pre-pregnancy body mass index (BMI) within overweight (19.9%) or obese (11.4%) categories¹³.

This report reviews: i) the association of maternal overweight and obesity with both adverse pregnancy outcomes and long-term adverse outcomes for mothers and babies; and ii) approaches that can be taken within primary obstetrical care to promote healthy weights among women who are either planning pregnancy, pregnant, or postpartum.

V. Overweight and Obesity Related to Childbearing

DEFINITIONS

Canadian Guidelines for the management and prevention of obesity in adults and children recommend measurement of both body mass index (BMI) and waist circumference to assess the level and distribution of adiposity in adults¹⁴. BMI is calculated as weight in kilograms divided by height in metres squared. Health Canada defines underweight as a BMI less than 18.5 kg/m², normal weight as a BMI of 18.5-24.9 kg/m², overweight as a BMI of 25.0-29.9 kg/m², class I obesity as a BMI of 30.0 - 34.9 kg/m², class II obesity as a BMI of 35.0 -39.9 kg/m², and class III obesity as a BMI \geq 40.0 kg/m^{2,15}. The other accepted measure of excessive body fat is waist circumference, based on the association between this measure of central adiposity and an increased risk of morbidity and mortality^{16,17,18}.

Waist circumference has replaced the waist-to-hip ratio since it is easier to perform. In adults with a BMI of 25 to 34.9 kg/m² (i.e. overweight or obese), a waist circumference greater than 88 cm (35 in) for women is associated with a greater risk of hypertension, type 2 diabetes, dyslipidemia, and coronary heart disease¹⁶.

There is ethnic variability in the BMI and waist circumference values that are associated with increased risk of adverse cardiovascular outcomes. The most appropriate cut-off values continue to be debated. The definitions discussed above apply to Caucasian, Hispanic and Black women. Lower cut-off values have been proposed for Asian women who have proportionately more truncal and visceral adiposity. BMI values of >23 kg/m² (overweight) and >30 kg/m² (obese), and waist circumferences of >80 cm have been proposed to define overweight among Asian women¹⁹.

These definitions of overweight and obese have not been rigorously applied in the relevant pregnancy literature. Definitions have included arbitrary weight cut-offs (e.g. obesity defined as ≥ 90 kg and normal weight as <75 kg²⁰). But when accepted thresholds have been applied, pre-pregnancy BMI categories have been used. The association between increased pre-pregnancy waist circumference and pregnancy complications has not been studied.

Maternal Weight

RISKS OF OVERWEIGHT AND OBESITY RELATED TO CHILDBEARING

A Medline search was performed from January 1980 to March 2008 using the following search terms: {'overweight' OR 'obesity' OR 'BMI' OR 'body mass index' OR 'weight gain'} AND {'pregnancy' OR 'pre-pregnancy'} AND {'risks' OR 'effects' OR 'complications' OR 'pregnancy outcomes'}. This search was limited to English language articles involving human subjects. We identified 2,993 articles of which 28 were included in this review, based on the following inclusion criteria: maternal weight was defined using BMI (rather than weight in kg), maternal and fetal outcomes were reported separately for overweight and obese categories, and measures of risk of overweight and obesity were relative to women of normal BMI.

Definitions of overweight and obesity in the included studies varied slightly but in general conformed to those which are currently accepted. There were 28 original studies^{7,9,21,22,23,24,25,26,27, 28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44}.

Fertility

Overweight and obesity have negative effects on fertility. Ovulatory disorders are more prevalent and frank anovulatory infertility increases in prevalence with increasing BMI⁴⁵. A common cause of anovulatory infertility is polycystic ovarian syndrome (PCOS)⁴⁶; obese women with PCOS have lower success rates with ovulation induction by pharmacological means⁴⁷.

Women who are overweight and obese have lower success rates with assisted reproduction. Specifically, being overweight or obese is associated in some (but not all studies) with longer periods of ovarian stimulation, higher doses of gonadotropins, lower oocyte retrieval, poorer oocyte quality, a lower incidence of embryo transfer, and lower rates of pregnancy^{48,49,50,51}.

During Pregnancy

Maternal

The risk of experiencing many complications in pregnancy increases as BMI increases above 24.9kg/m². However, the risk of experiencing some complications increases only as BMI increases above 29.9kg/m² (i.e. into the obese categories). In other words, women who are obese are at higher risk of experiencing complications than women who are overweight, but women who are overweight are still at higher risk than women of normal pre-pregnancy weight.

Gestational Diabetes (GDM)

Large observation studies have identified that the risk of gestational diabetes (GDM) is higher among overweight and obese women than women with a normal BMI^{7,8,9,21,25,28,29,30}. The magnitude of this risk increases progressively with increasing BMIs above 24.9. The association remains significant after adjusting for confounders such as maternal age, race, marital status, education, and parity⁵².

In a meta-analysis of 20 cohort studies⁵³, the unadjusted odds ratio (OR) for GDM was 2.14 (95% CI 1.82-2.53) for women who were overweight, 3.56 (95% CI 3.05-4.21) for class I obese, and 8.56 (95% CI 5.07-16.04) for class II/III obese, compared with women of normal weight pre-pregnancy. The associations were similar, albeit slightly lower, in a

Maternal Weight

separate meta-analysis which adjusted for confounders: OR of 1.86, 3.34 and 5.77 for GDM among overweight, class I obese, and class II/III obese women, respectively. British Columbian data are consistent with these findings. In 2005/6, the GDM rate was 6.8% of all pregnancies; however, 8.7% of overweight women and 14.4% of obese women had GDM¹³.

GDM is a pregnancy complication of importance because it is independently associated with a variety of complications for the mother, fetus, and baby in the index pregnancy^{54,55,56,57}, an increased risk of GDM in subsequent pregnancies, and an increased risk of type 2 diabetes for the mother and child^{21,55,56,58,59,60}. Estimates of the absolute risk of type 2 diabetes long-term have varied depending on the duration of follow-up and diagnostic criteria used. Fifty percent of women with a history of GDM will have developed diabetes after 10 years⁵⁸.

Hypertensive Disorders of Pregnancy (HDP)

As with GDM, the HDP are more common among women who are overweight and obese, with the risk increasing with increasing BMI. This excess of HDP includes both pre-existing hypertension and pre-eclampsia³⁶.

Pre-existing hypertension is more common among women who are overweight (OR 2.60; 95% CI 1.49-4.55) or obese (OR 7.93; 95% CI 4.74-13.27) compared with women of normal pre-pregnancy BMI (BMI 18.5-25kg/m²) (2,827 women)²⁵. Antenatal data from British Columbia showed that in 2005/6, hypertension occurred in 2.3% of women overall, but among 3.1% of overweight women and 5.8% of obese women¹³. These figures are similar to large studies from Europe (e.g. 3% and 5.3% for overweight and obese women, respectively; 25,601 women)³⁶.

Non-proteinuric gestational hypertension has been less well studied, but two recent cohort studies involving 24,241 women estimated that gestational hypertension risk was increased among overweight (OR 1.5, 95% CI 1.4-1.7), obese (OR 2.2, 95% CI 2.1-2.6), and class II/III obese women (OR 3.1, 95% CI 2.0-4.3)^{28,31}. It developed in 19.7%, 28.5%, 37.2% and 42.2% of normal weight, overweight, obese and class II/III obese women respectively³¹.

Pre-eclampsia risk increases among women who are overweight (OR 1.4- 2.3) or obese (OR 2.1-5.6)^{7,9,20,22,21,61,62}. Systematic review of observational literature (13 cohort studies, almost 1.4 million women) has estimated that pre-eclampsia risk doubles with each 5-to-7 kg/m² increase in pre-pregnancy BMI, even when women with confounding medical conditions (e.g. pre-existing hypertension or diabetes mellitus) or multiple gestations were excluded⁶³. In British Columbia (2005/6 data), the HDP are present among 7.3% of overweight and 11.6% of obese women, compared with 5.3% of women overall¹³.

Thromboembolism

Obesity is a well-recognized risk factor for thromboembolism^{20,64,65,66}, which is a leading cause of maternal mortality in Canada and worldwide⁶⁷. However, most studies addressing maternal weight-related pregnancy complications^{9,21,22,25,26,29} have not included thromboembolism as an outcome. When studied, risk increased with obesity (387 women, 129 cases and 258 controls⁶⁶, but also progressively with increasing weight. For example, the risk of antenatal thromboembolism doubled for women weighing 90-120 kg

Maternal Weight

and quadrupled for those who weighed more than 120kg in a Canadian population-based cohort study (142,404 women)²⁰.

Labour

Higher BMI is associated with prolonged gestation at term. In a study of 9,336 births, delivery at 42 weeks or more was more common among women who were overweight (OR 1.5; 95% CI 1.2-2.0) or obese (OR 1.7; 95% CI 1.2-2.3), compared with women of normal pre-pregnancy BMI³⁴.

Labour induction is more common among women who are overweight (OR 1.2-2.1) or obese (OR 1.6-3.2)^{7,22,25,28,31}. Failed labour induction is also more common^{28,68}. In a study of 126,080 deliveries, after excluding women with diabetes or HDP, obesity was associated with more failure to progress in the first stage relative to women with normal BMI (OR 3.1; 95% CI 2.5-3.8)⁶⁹. Labour duration is not affected by maternal weight²⁵.

Operative vaginal deliveries, third and fourth degree lacerations and shoulder dystocia are more common with increasing maternal BMI⁶⁸. Overweight and obese women are more prone to infections such as endometritis, urinary tract infections, wound infections and post operative pneumonia⁷. In one of the largest observational studies addressing pregnancy complications⁷, the risk of wound infection was increased among women who were overweight (OR 1.3; 95% CI 1.1-1.5) or obese (OR 2.2; 95% CI 1.9-2.6), compared with women of normal pre-pregnancy BMI. Similar figures were seen for urinary tract infection for overweight (OR 1.2, 95% CI 1.04-1.3) and obese women (OR 1.4, 95% CI 1.2-1.6), and genital tract infection among overweight (OR 1.24; 95% CI 1.09-1.41) and obese (OR 1.30; 95% CI 1.07-1.56) women.

Caesarean Section

BMI is positively associated with rates of both elective and emergency Caesarean delivery^{7,70,71,72,73} and inversely associated with the success of vaginal birth after prior caesarean delivery (VBAC)⁷⁴.

In a meta-analysis of 33 cohort studies involving 995,069 women, Caesarean delivery was more common among women who were overweight (OR 1.5; 95% CI 1.3-1.6), obese (OR 2.1; 95% CI 1.9-2.3), or who had class II/III obesity (OR 2.9; 95% CI 2.3-3.8)⁷⁵. Similar increases have been reported in other studies that adjusted for potential confounders such as pre-eclampsia, diabetes, and macrosomia^{21,28,29,30}.

Successful VBAC is inversely related to maternal weight. In a study of 510 women attempting a trial of labour, women who were underweight (BMI < 19.8 kg/m²) had the highest success rate (85%), compared with women who were overweight [from between pregnancies (74%) or from prior to the first pregnancy (71%)] and those women who were obese [from between pregnancies (56%) or from prior to the first pregnancy (55%)]⁷⁴.

The frequency of caesarean delivery among women who are overweight and obese is of concern due to increased rates of postoperative wound infection, endometritis, excessive blood loss, and thromboembolism^{7,21,70,76,77,78}.

Maternal Weight

Anaesthetic Complications

Anaesthetic risks among women who are overweight or obese are significant^{20,79}. The need for emergency anaesthesia is particularly hazardous. An anaesthetic review prior to labour should be considered. Women with class II/III obesity have the highest rate of complications, which include poor peripheral access, difficult placement of epidural or spinal anaesthesia, difficult intubation, more frequent pulmonary aspiration during anaesthesia, and more frequent sleep apnoea post partum (adjusted OR 2.0, 95% CI 1.3-3.1)²⁰.

Postpartum Haemorrhage

Studies have reported increased risk of post partum haemorrhage for women who are overweight (OR 1.1- 1.6) or obese (OR 1.2-1.7)^{7,25,28,31}, or who have class II/III obesity (OR 3.14)²⁸, compared with women with normal BMIs. However, the literature on this outcome is conflicting⁸⁰ possibly because the measurement of blood loss is subjective and the definition of postpartum haemorrhage variable. Women who are obese are at increased risk for post partum anaemia⁸¹ although no difference in the rates of blood transfusion has been demonstrated²⁰.

Breastfeeding

Several epidemiological studies have found an inverse association between maternal overweight and obesity and breastfeeding^{35,82,83,84,85}. In a large retrospective study, the likelihood of breastfeeding at discharge was significantly lower among mothers who were overweight (OR 0.86; 95% CI 0.84-0.88) or obese (0.58; 95% CI 0.56-0.60) compared to those with normal BMIs⁷. A systematic review of 22 observational studies found that after adjusting for confounders, women who were overweight or obese planned to breastfeed for a shorter period, were less likely to initiate breastfeeding, had delayed lactogenesis, and breastfed for shorter durations compared with women with normal BMIs⁸⁶.

Fetal and Perinatal Issues

Miscarriage

Women who are overweight or obese have an increased risk of miscarriage, following either natural conception or assisted reproduction^{49,87,88,89}. Following natural conception, women who are obese have a recurrent miscarriage rate that is more than three times higher than that for women of normal weight (OR 3.5, 95% CI 1.1-12.0)⁸⁸. A meta-analysis of 12 observational studies showed that women who were overweight or obese (vs. women of normal weight) were also more likely to miscarry (OR 1.3, 95% CI: 1.1-1.7) following in vitro fertilization (IVF).

Perinatal Mortality

Although between-study comparisons are hampered by different definitions of perinatal mortality, there appears to be an association between increasing maternal BMI and higher

Maternal Weight

perinatal mortality rates. For example, a Finnish study involving 25,601 singleton pregnancies demonstrated that perinatal mortality was higher among overweight (OR 1.5, 95% CI 1.0-2.4) or obese women (OR 2.2, 95% CI 1.3-3.6) compared with those of normal BMI³⁶. The association is strongest for stillbirth.

Stillbirth is more common among women who are overweight (OR 1.5, 95% CI 1.1-1.9) or obese (OR 2.1, 95% CI 1.6-2.7), compared with women with normal BMI (meta-analysis of 9 observational studies of primarily nulliparous women)⁹⁰. Although risk estimates were unadjusted, several cohort studies have demonstrated a significant positive association between maternal obesity and stillbirth after adjusting for factors such as maternal age, parity, diabetes, hypertensive disorders, smoking, alcohol, and chronic disease^{36,37,38,91}.

Neonatal death is more common among obese women. In a Danish cohort study involving 24,505 women with singleton pregnancies, obesity (vs. normal BMI) was associated with more neonatal death (adjusted OR 2.7; 95% CI 1.2-6.1)³⁸; similar rates were seen in a second study involving women with class II/III obesity³². In neither study was maternal overweight associated with neonatal death.

Congenital Malformations

There is a strong positive association between pre-pregnancy BMI and birth defects, particularly those involving the neural tube^{23,92,93,94,95,96}. A case-control study using data from the Atlanta Birth Defects Risk Factor Surveillance Study²³ demonstrated a 7% increase in the risk of fetal anomaly for each 1-unit incremental increase in BMI above a value of 25 kg/m² (i.e. any degree of overweight or obesity).

The National Birth Defects Prevention Study (NBDPS), the largest population-based study to date (with 10,249 cases and 4,065 controls) demonstrated an association between pre-pregnancy BMI and 16 categories of structural birth defects³⁹. Excluded from analyses were mothers with important confounders (e.g., pre-gestational diabetes) and adjustment was made for preconceptual folic acid consumption.

In the NBDPS, maternal obesity was most strongly associated with spina bifida (OR 2.1, 95% CI 1.6-2.7), as well as more severe forms of neural tube defect (i.e., anencephaly and hydrocephaly)^{92,94,96,97,98}. These data are consistent with older studies^{92,93,94,96,97,99,100}. No relationship has been demonstrated between maternal overweight and neural tube defects.

In the NBDPS study, maternal obesity was also positively associated with heart defects (OR 1.4, 95% CI 1.2-1.6) and diaphragmatic hernia (OR 1.4, 95% CI 1.03-2.0), and inversely associated with gastroschisis (adjusted OR 0.2; 95% CI 0.1-0.3). Maternal overweight was positively associated with heart defects (OR 1.1, 95% CI 1.01-1.3), omphaloceles (OR 1.5, 95% CI 1.-4-2.2), and hypospadias (OR 1.3, 95% CI 1.01-1.5).

The reasons for an association between maternal obesity and a spectrum of structural birth defects of different pathogenesis are unknown. For example, maternal overweight is associated with a higher risk of omphaloceles (which result from a failure of reduction of mid-gut herniation), whereas obesity is associated with more gastroschisis (which results from a vascular event). It must also be recognized that some associations have reached only borderline statistical significance and may be spurious. Regardless, undiagnosed maternal diabetes has been suggested in the pathogenesis of maternal weight-related fetal anomalies³⁹. Also, although folic acid supplementation was adjusted

Maternal Weight

for in the aforementioned studies, there is some evidence that obese women do not experience the typical reduction in neural tube defect risk associated with folic acid supplementation at the recommended 400 mcg/day⁹³.

Pre-term Delivery

The association between maternal BMI and pre-term delivery remains controversial with some studies showing an increased risk^{21,22,28,30,31,32,33} and others demonstrating a reduced risk or no change^{20,26,69}. When subtypes of pre-term birth are examined, however, infants of women who are overweight or obese have a higher risk of iatrogenic preterm delivery because of maternal complications, such as preeclampsia. The risk of preterm delivery due to spontaneous preterm labour appears to be reduced. This was demonstrated in a recent study which reported an increase in elective pre-term deliveries for women who were overweight (OR 1.2, 95% CI 1.03-1.3), obese (OR 1.5, 95% CI 1.3-1.8) or who had class II or III obesity (OR 2.1, 95% CI 1.8-2.6), but a lower risk for spontaneous preterm labour for women who were overweight (OR 0.9, 95% CI 0.8-0.98), class I obese (OR 0.9, 95% CI 0.7-0.99), and class II/III obese (OR 0.8, 95% CI 0.6-1.03)³². A large Danish study reported similar findings but in addition, noted a higher risk of preterm birth due to pre-term, premature rupture of membranes (PPROM) which has been associated with infections in the urogenital region³³.

Neonatal Morbidity

Infants of women who are obese are more likely to have low APGAR scores, hypoglycaemia, to require resuscitation, and to be admitted to a neonatal intensive care unit. In a retrospective cohort study, after adjusting for maternal age, smoking, parity and preexisting diabetes, the risk of an APGAR score ≤ 3 at five minutes was significantly higher for women who were overweight (OR 1.7; 95% CI 1.3-2.7), class I/II obese (OR 3.2; 95% CI 2.1-4.8), or class III obese (OR 6.0; 95% CI 2.7-13.4), compared with women of normal BMI²⁸. An Australian study reported that neonatal resuscitation was more common among women who were overweight (OR 1.3, 95% CI 1.0-1.7) or obese (OR 1.8; 95% CI 1.3-2.4), compared with infants born to women of normal weight²⁵. In addition, women who were obese were twice as likely to deliver an infant with neonatal hypoglycaemia (OR 1.9; 95% CI 1.2-3.1)²⁵. In a Canadian cohort study involving 18,643 women, the risk of NICU admission was higher among infants born to women who were overweight (OR 1.2; 95% CI 1.1-1.4), class I/II obese (OR 1.6; 95% CI 1.4-1.9), or class III obese (OR 2.9; 95% CI 1.9-4.4)²⁸.

Infant Birthweight

Maternal BMI has an important independent influence on infant birth weight. Several studies have demonstrated a protective effect of maternal overweight on delivery of small for gestational age (SGA) infants^{26,30,36}. Macrosomia or a large for gestational age (LGA) infant are also more common among women who are overweight (OR 1.3- 2.1) or obese; this effect persists after adjustment for maternal diabetes (OR 1.9-3.5)^{7,21,22,28,29,30,31}. Macrosomia is a powerful predictor of shoulder dystocia and through this, third and fourth degree perineal lacerations, blood loss, and neurological injury to the infant^{27,101}. Macrosomic infants face an increased obesity later in life¹⁰².

Maternal Weight

The Risks in Perspective

It is clear that there are substantial pregnancy-associated risks of a BMI above 25 kg/m². The attributable risks are also high given the prevalence of overweight and obesity in the general obstetric population (i.e. 30% in British Columbia)¹³.

The pregnancy-related risks of obesity are comparable in magnitude to other well-recognized antenatal risk factors. For example, the risk of preterm delivery at <33 weeks among women who are obese (OR 1.5 - 2.0 compared with women of normal weight)^{21,30,31} is equivalent to that associated with maternal smoking of up to 10 cigarettes daily (OR 1.4-1.5 compared with non-smokers)^{30,103}. An alcohol intake of more than 5 standard drinks per day triples the risk of stillbirth (OR 3.65 compared with non-drinkers)¹⁰⁴, as does a maternal BMI over 30 kg/m²^(26,91). Neural tube defects are three times more common among women who are obese compared with those with a normal BMI²³. This amounts to an absolute risk increase of about 1%, similar to early pregnancy exposure to carbamazepine or folic acid antagonists (OR 2.8)¹⁰⁵. As such, reducing women's weight pre-pregnancy should be given similar priority to that of maternal smoking cessation, alcohol cessation, adequate folic acid supplementation, and avoidance of teratogenic drugs.

MEDICAL ASSESSMENT CONSIDERATIONS/ CHALLENGES

Several practical difficulties can arise in managing care for women who are overweight or obese. Blood pressure is overestimated when standard cuffs are used¹⁰⁶. Abdominal palpation to assess fetal growth, lie or presentation may be difficult due to increased adiposity. Prenatal sonographic diagnosis is difficult, particularly if BMI is >36kg/m²⁽¹⁰⁷⁾. Ultrasound visualization of fetal anatomy declines with increasing maternal size, with the most marked reductions in visualization for fetal heart, umbilical cord, and spine¹⁰⁷. Ten percent of women who are obese have suboptimal 4-chamber views at 22-24 weeks gestation despite the use of advanced ultrasound equipment¹⁰⁸. Fetal monitoring by intermittent auscultation or continuous cardiotocography (CTG) using external abdominal transducers can be technically difficult. The depth of maternal adipose tissue can interfere with the doppler signal which may compromise CTG quality and contribute to poor neonatal outcomes¹⁰⁹.

LONG-TERM MATERNAL AND PAEDIATRIC IMPLICATIONS

Obesity is associated with increased all-cause mortality for women^{110, 111,112,113}. The risks for a number of health conditions increase with increasing BMI. Several factors related to cardiovascular disease increase with increasing BMI¹¹⁴, including stroke¹¹⁰, coronary heart disease^{110,112}, hypertension^{110,112}, and hyperlipidemia¹¹². Type 2 diabetes increases with increasing BMI^{110,112}, as does kidney disease^{115,116}. The risk for a number of cancers increases, including endometrial cancer, breast cancer, gallbladder cancer, oesophageal cancer, renal cancer, leukemia, multiple myeloma, and non-Hodgkin lymphoma^{110,117}. Rates of asthma¹¹⁸ and arthritis^{119,120} are higher among women who are obese. The risk of dementia may increase with increasing BMI¹²¹.

Maternal obesity, both independently and through association with diabetes, is a risk factor for childhood obesity¹²² and obesity in early adult life^{57,123,124,125,126,127,128,129,130,131}. Children of women who were obese during early pregnancy are twice as likely to be obese at ages 2, 3 and 4¹³², and almost 3 times as likely at age 7¹²² compared with children whose mothers had a normal pre-pregnancy BMI. A dose-response relationship was

Maternal Weight

shown in a longitudinal study of 6,280 participants in which increasing maternal BMI was associated with increasing BMI of the offspring at ages 1, 14 and 31 years¹²³.

These effects may be due to more than genetics or postnatal environment¹³³. Fetal programming may occur whereby *in utero* exposure to overnutrition (secondary to overweight/obesity) and/or hyperinsulinaemia (secondary to maternal diabetes), results in long-lasting changes to fetal metabolism and this 'thrifty' (i.e. energy-saving) phenotype increases the risk of obesity or diabetes in later life^{57,134}. For example, maternal diabetes increases the offspring's risk of type 2 diabetes more than would be predicted from genetics alone¹³³. In a recent American study involving nearly 10,000 multiethnic mother-child pairs, increasing degrees of hyperglycaemia in pregnancy were associated with higher rates of obesity in their children at age 5-7 years¹³⁵. Importantly, treatment of GDM attenuated these risks¹³⁵.

COST IMPLICATIONS

There has been limited research into the cost implications of overweight and obesity as related to childbearing. During pregnancy, two studies from France have shown that the cost of prenatal care is 5.4- to 16.2-fold higher for women with a BMI of 25-34 kg/m², compared with that for women with normal BMI^{136,137}. When both prenatal and postnatal care were considered, cost was even higher, due in large part to more prolonged hospitalization (by an average of 4.4 days), compared with women of normal BMI. In an Australian obstetric population, prolonged hospitalization (exceeding 5 days) was also more common among women who were overweight (OR 1.4, 95% CI 1.1-1.6), class I/II obese (OR 1.5, 95% CI 1.2-1.9), and class III obese (OR 3.2, 95% CI 2.2-4.6), compared with women of normal weight⁹. Hospitalization was also longer in an American study of 13,442 pregnancies; after adjustment for age, ethnicity, level of education and parity, the mean (+/-SE) hospital stay for delivery was greater among women who were overweight (3.7±0.1 days), obese (4.0±0.1 days), class II obese (4.1±0.1 days), or class III obese (4.4±0.1 days), compared with women of normal BMI (3.6±0.1 days). Most of the increase in length of stay associated with higher BMI was related to increased rates of Caesarean delivery, maternal weight-related anaesthetic risks and postpartum complications¹³⁸.

Although longer maternal length of stay is a major contributor to cost, other contributors include: extra pregnancy screening at maternity clinics, anaesthetic risks, and postpartum complications for the mother, and costs relating to increased neonatal intensive care unit admission for the baby.

In the same American study quoted above¹³⁸, overweight and obesity were also associated with more prenatal fetal tests, obstetrical ultrasonographic examinations, medications dispensed from the outpatient pharmacy, telephone calls to the department of obstetrics and gynaecology, and prenatal visits with physicians, but fewer prenatal visits with nurse practitioners and physician assistants.

VI. Gestational Weight Gain

GUIDELINES

Weight gain in pregnancy is generally considered to be the difference between a woman's weight at the last antenatal visit and either her pre-gravid weight or her weight at the first antenatal visit.

Maternal Weight

The 1999 Canadian guidelines by Health Canada were adapted from the 1990 Institute of Medicine (IOM) guidelines¹³⁹ developed to reduce the risk of fetal growth restriction. Gestational weight gain goals are adjusted based on pre-pregnancy BMI. Table 1 presents the guidelines for singleton pregnancies; for women with multiple gestations, weight gain goals are slightly higher¹⁴⁰. According to these guidelines, healthy weight pregnant women with twins should gain a total of 16.0-20.5 kg (35-45 lb.) and aim for a rate of gain equal to 0.7 kg (1.5 lb.) each week during the second and third trimesters.

Several studies have confirmed that pregnancy weight gain within the IOM ranges optimizes term deliveries of babies of birthweight between 3,000 and 4,000 g^{141,142,143}. However, there is concern that IOM guidelines for gestational weight gain are too high, especially for women who are overweight or obese^{144,145}. The IOM is currently reviewing their guidelines with the revised guidelines expected in June 2009. Health Canada is observing the IOM review with the intention to revisit the Canadian guidelines.

RISKS OF EXCESSIVE GESTATIONAL WEIGHT GAIN

Gestational weight gain in excess of the guidelines is associated with an increase in both maternal complications (such as GDM⁶⁸), pre-eclampsia^{146,147}, failed induction of labour, or Caesarean section^{68,148,149}, and perinatal complications (such as fetal macrosomia or LGA infants and fetal distress^{142,146,147,148,149,150,151}), after adjusting for confounders such as maternal age, race, parity, and smoking. The magnitude of increased risk is smaller than that seen among women who are either overweight or obese pre-pregnancy, with OR of about 1.5 for all but LGA infants with an OR of over 2. As such, one sees a dose-response relationship between maternal weight and adverse pregnancy outcomes, such that risk is: increased with excessive gestational weight gain; greater with pre-pregnancy overweight; and greatest with pre-pregnancy obesity. As with maternal overweight and obesity, excessive gestational weight gain is also a risk factor for childhood obesity¹⁵².

Pregnancy weight gain in excess of the IOM guidelines is common. The range of weight gain among women with normal pregnancy outcome is so variable that gestational weight gain *alone* is not a good predictor of pregnancy outcome¹⁵³. Although obese women tend to gain less weight overall^{154,155,156,157}, they are more likely to gain weight in excess of guidelines, compared with women of normal weight¹⁵⁸. In British Columbia (2005/6 data), the mean weight gain (\pm SD) in pregnancy was similar among women of normal weight (15.7 \pm 5.5 kg), overweight (14.8 \pm 6.4 kg), and obese (11.7 \pm 7.2 kg). However, women of normal pre-pregnancy weight were less likely to have excessive weight gain (39.3%) compared with women who were overweight (60.4%) or obese (65.7%)¹³.

Risk markers for excessive weight gain include: younger age, being single, Black, or Hispanic¹⁵⁹, or self-report of low numbers of supportive individuals in women's lives¹⁶⁰. Excessive weight gain is associated with either a lack of provider advice about recommended weight gain, or, advice to gain more than the IOM guidelines for women who are overweight or obese¹⁶¹.

All of the following dietary patterns have been associated with excessive weight gain, particularly for women who are overweight pre-pregnancy: a diet that is high in energy¹⁶², fat or glycaemia^{163,164}, and consumption of sweets¹⁶¹. However, a change in eating habits during pregnancy may be more closely related to weight gain than the actual macronutrient ratio of the foods chosen¹⁶².

Maternal Weight

Pre-pregnancy dieting or restrained eating are also risk factors for excessive gestational weight gain^{165,166}, possibly because they may impair an individual's ability to respond to hunger and fullness cues¹⁶⁶. Pre-pregnancy dieting is of particular concern because it is so common. In the 1999 BC Nutrition Survey, 46.4% of women aged 19 – 30 and 42.5% of women aged 31-50 reported trying to lose weight; 72.9% of them by changing their eating habits¹². During pregnancy, women with a history of dieting report the perception of being “allowed” to eat more food, and “allowed” to eat “fattening” foods that were previously perceived to be off-limits¹⁶⁵.

VII. Supporting Women and Health Care Providers to Reduce Risks

The 2003-2005 Confidential Enquiry into Maternal and Child Health (CEMACH) included in their list of top 10 recommendations the urgent development of national guidelines for management of obese pregnant women in the United Kingdom. A key finding of the Enquiry was that more than half of all the women who died from direct or indirect causes (and for whom information was available) were overweight or obese, and more than 15% were class III obese⁶⁷. There are no published guidelines on managing pregnancy among overweight and obese women from the Society of Obstetricians and Gynaecologists of Canada (SOGC).

While an increased risk of many complications has been identified in association with overweight or obesity during pregnancy, there is insufficient evidence to identify best practices to reduce these risks.

INCORPORATE BEHAVIOUR CHANGE THEORY & PRACTICES

Women and their behaviors do not exist within a vacuum. The contexts of women's lives provide both motivations and barriers to adopting behavior change. As stated in the World Health Organization's Ottawa Charter for Health Promotion: “health is created and lived by people within the settings of their everyday life; where they learn, work, play and love”¹⁶⁷. Therefore, women-centred practice is recommended to promote health¹⁶⁷. A women-centred approach to maternity care is one in which services are located in the context of primary care with the recognition that for the majority of women, pregnancy and childbirth are normal life events¹⁶⁸. In this approach, maternity services place the mother and her baby at the centre of care, and plan and provide services to meet their needs¹⁶⁸. Developing women-centred care relies, in part, on understanding women's preferences and needs with respect to care¹⁶⁸. It also involves engaging women and their families (as defined by the woman) in the processes of planning and delivering services¹⁶⁸.

Behaviour change requires both recognition of the problem to be addressed, and adoption of new behaviours. Motivational interviewing is emerging as a promising practice for promoting healthy weight among several populations; however, it has not yet been established among women during the perinatal period^{169,170,171,172,173,174,175,176}.

A foundational tenet of motivational interviewing is that people vary in their readiness for change and can be influenced to increase their motivation for making changes¹⁷⁷. Motivational interviewing is often used along with Prochaska and DiClemente's transtheoretical model of change (also known as the stages of change theory¹⁷⁷). While the specific delineation of the stages has been questioned, the transtheoretical model of

Maternal Weight

change identifies that some people are not even thinking about changing a behaviour (known as precontemplation) while others are thinking about and taking steps towards making a behaviour change¹⁶⁹. The model states that practitioners should use different techniques to assist individuals with making a behaviour change depending on whether they are precontemplators or women thinking about/taking steps towards change¹⁷⁸.

CHECKLIST

Table II outlines a checklist of reasonable practices to support women who are overweight or obese (in order to improve maternal and perinatal outcomes) and to support practitioners to initiate appropriate consultations and investigations; it has been modified from two sources^{179,180}. The checklist includes issues to address when women are planning pregnancy, are pregnant, or are postpartum.

PRE-PREGNANCY

Healthy Eating, Physical Activity, Substance Reduction and Avoidance

Health Canada recommends that all women of childbearing age follow the healthy eating recommendations in Canada's Food Guide¹⁸¹ and be physically active every day following Canada's Physical Activity Guide¹⁸².

To promote tobacco reduction and cessation, the British Columbia Perinatal Health Program Guideline: Tobacco Use in the Perinatal Period recommends that physicians: ask women of childbearing age about their smoking status; advise those who smoke how important it is to stop and avoid exposure to second hand smoke; and, offer assistance in quitting to those who smoke¹⁸³.

To promote alcohol and other substance use reduction and cessation, the British Columbia Perinatal Health Program Guideline: Alcohol Use in the Perinatal Period and Fetal Alcohol Spectrum Disorder recommends that physicians: talk about substance use with all women; identify those who need help with reducing drinking and other substance use; and support women to reduce or stop use of alcohol and other substances¹⁸⁴.

Weight Loss

The American College of Obstetricians and Gynecologists (ACOG) recommends preconception counselling about potential complications of maternal overweight and obesity, and participation in weight loss programs pre-pregnancy¹⁸⁵.

Weight loss may improve fertility. A loss of 5-10% of body weight may restore ovulation within six months for more than half of obese women with polycystic ovarian syndrome (PCOS)^{186,187,188,189}. If conception is not yet desired, contraception should be advised, even for women with demonstrated subfertility. Several studies have shown that following bariatric surgery, fertility is improved¹⁹⁰.

It is assumed that weight reduction before pregnancy will improve future pregnancy outcomes, but there are no randomized controlled trials that have established that this assumption is correct. A cohort study involving 150,000 Swedish women with two births examined changes in BMI between pregnancies in relation to perinatal morbidity. Women with a BMI decrease of more than one unit had significantly lower rates of pre-

Maternal Weight

eclampsia and LGA infants in their second pregnancy. However, as only 5.4% of women in this study were obese and 19% were overweight, it is unclear whether such modest weight loss in overweight or obese women would result in similar benefits¹⁹¹. Several studies have shown that following bariatric surgery, adverse maternal (e.g. pre-eclampsia, GDM) and fetal outcomes (e.g. macrosomia) are reduced in a subsequent pregnancy¹⁹⁰. If a woman has undergone bariatric surgery prior to conception, she may need supplementation with vitamin B12, iron, calcium, and folic acid^{192,193,194}.

Regardless of proven benefit on subsequent pregnancy outcomes, there are recognized health benefits for women for moderate weight reduction (on glycaemic control, BP, lipids, and the risk of stroke, coronary artery disease and death) that make a convincing argument for advocating for taking steps towards the achievement of a healthy weight outside of pregnancy.

Outside of pregnancy, better practices for the achievement of a healthy weight involve lifestyle interventions including calorie reduction, physical activity, and behavioural therapy. Pharmacotherapy and bariatric surgery may also have a role. Canadian clinical practice guidelines recommend consideration of pharmacotherapy for individuals with BMI ≥ 27 kg/m² in the presence of cardiovascular risk factors or BMI ≥ 30 kg/m² if 3-6 months of lifestyle modification has been unsuccessful. Bariatric surgery could be considered for individuals with BMI ≥ 35 kg/m² and cardiovascular risk factors, or BMI ≥ 40 kg/m² if other weight loss strategies have failed¹⁴.

There are potential barriers to achieving weight loss among women who are overweight or obese and planning a pregnancy. First, 50% of pregnancies are unplanned. Second, as the most successfully maintained weight loss involves lifestyle change resulting in slow progressive weight loss, it is possible that women who are planning pregnancy may not be willing to wait for such a long time prior to conceiving. In the setting of advanced maternal age, it may even be *inappropriate* to suggest that conception be delayed to achieve weight loss. There appears to be a consensus, however, that pregnancy should be delayed for two years following bariatric surgery, during which time one sees the majority of the weight loss and postoperative complications.

As we believe that either the postpartum period or the period before women are planning a pregnancy may be the times to target weight loss, barriers to success and strategies to achieve weight loss are discussed under '*Supporting Women and Health Care Providers to Reduce Risks/ Postpartum*'.

Folic Acid Supplementation

For all women of childbearing age (including during pregnancy), a multivitamin and mineral supplement containing 0.4 mg daily of folic acid is recommended to reduce the risk of NTDs and possibly, other congenital anomalies¹⁸¹. Women with a BMI > 30 kg/m² may need higher doses of folic acid to achieve the same folate levels as women with a BMI < 20 kg/m²⁽¹⁹⁵⁾. Although the mechanisms for the obesity-associated risk of NTDs are poorly understood, the 2007 guidelines of the Society of Obstetricians and Gynaecologists of Canada (SOGC) recommend that women with class II/III obesity take 5 mg daily of folic acid for at least three months before conception, and continue this dose until 10-12 weeks post-conception¹⁹⁵.

Maternal Weight

Optimize Treatment of Co-morbidities

Chronic diseases such as type 2 diabetes, hypertension, and obstructive sleep apnea are more common among women who are overweight or obese¹⁹⁶. Treatment of these co-morbidities should be optimized prior to conception. In addition, early assessment of glycaemic control, liver enzymes, renal function, and proteinuria should be considered as this may help to later distinguish pre-pregnancy morbidity (such as type 2 diabetes, macrohepatic steatosis, or pre-existing renal disease) from pregnancy-induced complications [such as gestational diabetes, HELLP syndrome (hemolysis, elevated liver enzymes, low platelets), or gestational proteinuria of pre-eclampsia]. Both pre-pregnancy morbidity and pregnancy complications occur more commonly in association with overweight and obesity¹⁹⁷.

DURING PREGNANCY

Multidisciplinary Support

Given the array of maternal and fetal complications that may occur for women who are overweight or obese, involvement of a multidisciplinary team should be considered, similar to that which is commonly available for women with diabetes.

Screen for GDM

Screen women for GDM following the British Columbia Perinatal Health Program Obstetric Guideline 10A: Gestational Diabetes¹⁹⁸. This guideline recommends that women with identified risk factors be screened in the first or second trimesters.

Set an Appropriate Weight Gain Goal

Height, weight and BMI should be recorded on the first antenatal visit. At present, height and weight are each recorded on 80% of antenatal records in British Columbia (2005/6 data), allowing for calculation of BMI in 70% of women¹³.

An appropriate gestational weight gain goal should be set based on current Canadian Guidelines (based on the IOM guidelines), and risks associated with excessive weight gain should be discussed in early pregnancy. Based on results of questionnaire-based studies, prenatal care providers are not following IOM guidelines when advising patients about gestational weight gain. In two recent studies^{199,200}, a third of women reported having received no advice from their health providers about how much weight to gain during pregnancy. And, among women who did receive advice, 50% of women who were overweight were advised to gain more than recommended by guidelines²⁰⁰.

Weight loss is not recommended during pregnancy for all women, including women who are overweight or obese, because of concerns about the impact of maternal ketosis on fetal brain development²⁰¹.

Maternal Weight

Encourage Healthy Eating and Physical Activity (to achieve weight gain goal)

Healthy Eating

There are limited data on actual nutritional intake because pregnant women are usually excluded from analyses. Available information suggests that women do not meet nutrition recommendations for pregnancy. American data from 1988-94 reported that women consumed insufficient grains, vegetables, fruit, milk and meat, and consumed excessive fat, saturated fat, and sodium²⁰². This finding is in contrast to the fact that women report wanting to eat well to support their fetus' health²⁰³, and appear to have adequate nutrition knowledge, regardless of socioeconomic status²⁰⁴.

Physical Activity

Physical activity during pregnancy is advisable, not just acceptable, for all pregnant women.

In small studies, physical activity in pregnancy was not associated with adverse maternal or perinatal outcomes^{205,206,207,208,209,210,211,212,213,214}. In fact, lack of physical activity is associated with loss of muscular and cardiovascular fitness, excessive maternal weight gain, higher risk of gestational diabetes, higher risk of pregnancy-induced hypertension, development of varicose veins and deep vein thrombosis, and more physical complaints such as dyspnea or low back pain²¹⁵.

Physical activity also has the potential to positively influence body image satisfaction²¹⁶. Among 74 Caucasian women of high socioeconomic status, the perceived benefits of exercise in pregnancy were (in order of importance): improved mood, increased energy/stamina, and assistance with fitness and weight control²¹⁷. Reported barriers to physical activity during pregnancy were feeling too big and physical limitations, such as nausea, vomiting and fatigue²¹¹.

The SOGC guideline (2003) recommends maintenance of a good fitness level throughout pregnancy without trying to reach peak fitness or train for an athletic competition²¹⁵. Suggested activities include brisk walking, stationary cycling, cross-country skiing and swimming as they are associated with minimal risk of loss of balance or fetal trauma.

Barriers to Healthy Eating and Physical Activity

Related to Health Care Providers

Conversations about healthy eating and activity during pregnancy may not be taking place, or incorrect information may be communicated²⁰³. As such, women may not know the recommended weight gain for their pre-pregnancy BMI and/or that it is both safe and advisable to exercise during pregnancy. Most studies suggest that women value having conversations with their health care providers about healthy weight, healthy eating, and physical activity^{199,164, 203,218}.

There is opportunity for health care providers to better communicate the physical activity recommendations for pregnancy, as many women believe that physical activity in

Maternal Weight

pregnancy is unsafe. Women may be discouraged by family and friends or be discouraged by what they perceive as mixed advice from health care providers²¹⁹.

In a survey of members of the American College of Obstetricians and Gynecologists, most members were aware of obesity-related pregnancy risks and felt that weight management fell within their clinical practice but did not feel adequately trained in weight management²¹⁸. We could not identify relevant Canadian data. Other barriers to weight management included both practice-related factors (e.g. support services, visit time) and patient-related factors (e.g. patient motivation and compliance with dietary and activity advice)²¹⁸. Of interest is the fact that when health care providers did not influence women's lifestyle, their spouse/ partner often did²¹⁷.

Related to Women

There are limited data on barriers to meeting nutrition recommendations in pregnancy. However, as discussed above (under '*Encourage healthy eating and physical activity*'), barriers to healthy eating do NOT appear to include lack of nutrition knowledge²⁰⁴.

A barrier to healthy lifestyle in pregnancy may be the interaction between how women feel about their bodies and their eating and physical activity habits, of which one investigation identified four patterns based on attitudes during but also before and after pregnancy²²⁰:

1. **Relaxed maintenance:** Women in this category are well educated. They have stable weight and are content with their weight pre-pregnancy. During pregnancy these women "watch" what they eat, eat a "healthy" diet, and are physically active in their daily lives (but do not intentionally plan exercise). They are not concerned about their weight during pregnancy. Postpartum weight is not of much concern and they return to a healthy eating pattern similar to their pre-pregnancy habits. They continue to include physical activity in daily life but do not plan exercise. Their postpartum weight returns to close to pre-pregnancy weight but they may maintain some extra weight.
2. **Exercise:** Women in this category are well educated. Pre-pregnancy exercise and fitness are priorities for these women. They have healthy eating habits pre-pregnancy and are happy with their weight. They maintain healthy eating during pregnancy and continue to include exercise while pregnant, but less frequently and intensely than pre-pregnancy. Exercisers return to exercising postpartum, some after a non-active transition time. Their motivation for loss of their pregnancy weight is related to both health and body image. They don't retain weight postpartum but those who take a non-active transition time take longer to return to their pre-pregnancy weight.
3. **Determined:** Women in this category have a variety of educational levels. They perceive themselves as overweight pre-pregnancy, but they may or may not have tried losing weight. These women are somewhat uncomfortable with gaining weight during pregnancy. They are determined to lose weight postpartum and achieve a lower weight than pre-pregnancy. Postpartum, they go on diets and exercise to lose weight. They are frustrated at the slow rate of weight lost postpartum but they do reach their pre-pregnancy weight by approximately one year.

Maternal Weight

4. **Unhurried:** These women have the lowest education. They are unhappy with their weight pre-pregnancy but most do not take action to lose weight. They do not plan physical activity during pregnancy and have unhealthy eating habits (e.g. consume more high-fat and high-sugar sweets and snacks than women in the other three groups). They are unhappy with their weight postpartum but do not make weight loss a priority and take no action (eating habits or physical activity).

In order to plan effective interventions, it is important to understand how women respond and adjust to the changes in eating, body weight/ shape, and physical activity that are associated with pregnancy and the period following delivery²²¹.

Effectiveness of Lifestyle Interventions

Lifestyle interventions during pregnancy are aimed at curbing gestational weight gain; they are NOT aimed at weight loss. Interventions aimed at curbing excessive gestational weight gain have had mixed results in a variety of settings and with women of variable socioeconomic status and pre-pregnancy weight^{222,223,224}.

No effect was seen on gestational weight gain, eating habits (except for caffeine intake), or physical activity levels following individual dietary counselling and physical activity sessions among aboriginal Cree women²²⁵. Gestational weight gain and physical activity were unaltered, although there were some dietary improvements (increased fibre, fruit and vegetable intake) following dietary and physical activity counselling from early pregnancy, provided by public health nurses at antenatal visits, for primiparous women in Finland²²⁴.

In contrast, excessive weight gain was curbed among women who were normal weight (but not overweight) by a behavioural intervention (vs. usual care) in a randomized controlled trial involving low-income women²¹⁶. Weight gain was curbed among normal and overweight low-income (but not high-income) women by providing education on weight gain, diet, and exercise by mail, as well as guidance about and monitoring of gestational weight gain by health care providers²¹⁷. Gestational weight gain was reduced by weekly motivational talks and regular physical activity among obese women in Sweden²²⁴; modes of delivery and neonatal outcomes were not changed.

The mixed results of existing intervention studies emphasize that there are no simple solutions to supporting women to curb excessive gestational weight gain. These data should serve to motivate the development of better interventions based on published successes and failures. For example, there may be a role for education and/or involvement of spouses and partners, because when health care providers did not influence women's lifestyle, their spouse/ partner often did²¹⁷. There may also be a role for novel approaches that take into account women's barriers to healthy lifestyle adoption, as well as their motivation (i.e. readiness for change). For example, one could consider focussing on women who demonstrate a 'determined' pattern of eating/ lifestyle, rather than on those who are 'unhurried' and who may, therefore, be less receptive to interventions.

Maternal Weight

POSTPARTUM

Prevention of Weight Retention

The pattern of postpartum weight loss, and the amount of weight ultimately retained postpartum, is variable. In general, women lose weight gradually, reach their lowest BMI (which usually includes some weight retention) at approximately 12 months^{226,227}, and then gain weight slowly thereafter^{160,226}.

Postpartum weight retention is of concern because of its potential impact on subsequent pregnancy weight gain²²⁸ and long-term overweight and obesity. Even if a woman's weight is higher than pre-pregnancy but remains within the normal range, there is an increased risk of excessive gestational weight gain and adverse pregnancy outcomes in a subsequent pregnancy¹⁹¹.

Postpartum weight retention is most strongly associated with excessive weight gain in pregnancy, particularly excessive first trimester weight gain^{52,154,159,160,227,229,230,231,232,233,234,235,236,237}. In fact, excessive weight gain in pregnancy may explain some of the association between postpartum weight retention and both overweight and obesity²³⁸. There is mixed evidence regarding whether postpartum weight retention is associated with pre-pregnancy overweight or obesity^{228,229,238,239}. It may be that pre-pregnancy overweight or obesity indirectly increases the risk of postpartum weight retention because women who are overweight or obese are at higher risk of gaining more than the IOM guidelines²³⁸. And, gaining more than the IOM guidelines, in turn, increases risk for postpartum weight retention²³⁸.

There are many other risk factors for postpartum weight retention, including: unhealthy eating habits²⁴⁰, low physical activity^{160,241}, inadequate sleep²⁴², low socioeconomic status/education^{229,240}, low affect/ depression²²⁹, ethnicity^{159,227}, and self-report of low numbers of supportive individuals in women's lives¹⁶⁰. In the United States, Black women retain more weight than Caucasian women, and both Black and Caucasian women retain more weight than Asian women^{159,227}. There is a lack of evidence regarding weight retention among women of other ethnic backgrounds.

As risk factors for postpartum weight retention, inadequate sleep and mood are worthy of specific mention because each is so prevalent, and because each is potentially modifiable. A study involving 940 Caucasian women with higher socio-economic status found that women who reported less than five hours of sleep per night at six months postpartum were more likely to retain 5kg or more at one year postpartum²⁴². Postpartum, low affect (or "the blues") occurs among 70-80% of women and frank depression in 12-16%²⁴³. It appears that overweight and obesity may cluster with depression and disordered eating (preoccupation with body shape and pathological avoidance of "fattening" foods) postpartum²⁴⁴.

Any effect of breastfeeding on postpartum weight retention is small, but there are, of course, many other reasons to breastfeed. Breastfeeding (especially exclusively without supplementing with formula) may promote a small amount of postpartum weight loss²⁴⁵, and women who choose to breastfeed have better diet quality^{238,246}.

Maternal Weight

Weight Loss Strategies

Outside Pregnancy

The Canadian Medical Association 2006 Canadian clinical practice guidelines on the management of obesity in adults and children recommend a comprehensive healthy lifestyle intervention for people who are overweight and obese that includes an energy-reduced diet, regular physical activity, and education and support in behaviour modification techniques¹⁴. Health care providers are also encouraged to create a non-judgemental atmosphere when discussing weight, and to consider an individual's barriers to weight loss¹⁴.

Among the adult population in general, the weight loss industry is large, and many lessons can be learned from it. The variety of approaches that have been tried is detailed under '*During pregnancy*'. Most interventions are still only moderately successful in promoting sustained, small amounts of weight loss²⁴⁷. The evidence is weak for determining best practice regarding weight management for overweight and obese individuals²⁴⁸ and there is insufficient evidence to determine the effectiveness of commercial weight loss programs²⁴⁹.

The following weight loss delivery methods have been identified as warranting future investigation:

- Reminder systems for physicians to perform specific actions²⁴⁸,
- Brief training interventions for health care providers²⁴⁸,
- Shared care between a hospital program and GP²⁴⁸,
- In-patient care, and dietitian-led treatments²⁴⁸,
- On-going monitoring and support²⁵⁰, and
- Long-term follow-up²⁵⁰ particularly personal contact (vs. internet-based follow-up or no follow-up at all)²⁵¹.

The following weight loss program content has shown promise:

- Setting realistic weight loss goals²⁵⁰,
- Teaching eating behaviour control techniques (cognitive-behaviour therapy)^{252,253},
- Promoting healthy eating *and* physical activity²⁵⁴,
- Teaching self-esteem²⁵⁵.

The evidence is weak regarding the optimal diet composition for weight loss and maintenance, especially considering high drop-out rates and poor diet adherence²⁵⁶. Low glycemic index diets may promote weight loss and weight maintenance^{257,258}. Lower carbohydrate diets may promote greater short-term weight loss without superior long-term effectiveness^{259,260}. As such, there may not be one 'optimal' diet for everyone. Considering the high drop-out rates, an 'optimal' diet may be an eating pattern that an individual is able to follow long-term and so there may be as many 'optimal' diets as there are individuals²⁵⁹.

Maternal Weight

Tailoring Weight Loss Programs to Women Who are Postpartum

The American College of Obstetricians and Gynecologists recommends participation in nutrition counselling and exercise programs postpartum¹⁸⁵. A Cochrane review (as well as the National Research Council and Institute of Medicine¹⁶¹) concluded that there is insufficient evidence to identify best practices for the promotion of postpartum weight loss and prevention of weight retention²⁶¹. Study drop-out rates are high and can exceed 40%.

A dietary component of postpartum weight loss programs appears to be important. In fact, diet and exercise are not more effective than diet alone²⁶¹. On average, postpartum weight loss interventions have resulted in a weight loss of 4.8-7.8kg²⁶². Greater weight loss is seen with programs of longer duration.

There is mixed evidence regarding whether women change their eating habits postpartum compared to pre-pregnancy^{238,263}. Even if they do improve their eating habits, women during the postpartum period do not consume the recommended amount of grain products, vegetables and fruit, or milk and alternatives^{238,263}.

Women who are overweight or obese report different eating patterns than women with normal BMIs. A study involving 340 low-income women found that those who were obese one year postpartum reported more barriers to healthy eating, more emotional eating, and more food insecurity (i.e. a lack of access to safe, affordable, healthy food) than women who had normal BMIs²⁴⁰. Women who had normal BMIs one year postpartum reported eating in response to taste, hunger, and cravings more than women who were overweight or obese²⁴⁰.

Physical activity alone has not been successful in achieving postpartum weight loss, although physical activity may have other benefits. Group physical activity may decrease postpartum depression, but it is not clear whether the physical activity or the opportunity to interact with other new mothers was responsible for the improved mood (2 RCTs, 39 women)²⁶⁴. Unfortunately, studies show that most women decrease strenuous, moderate, and mild physical activity during pregnancy and postpartum versus pre-pregnancy^{160,217,241}.

Being physically active postpartum is related to self-efficacy (i.e. believing that you can do it²⁶⁵), and being physically active pre-pregnancy²⁴¹. In a study involving 74 Caucasian, high socioeconomic status women who were physically active during pregnancy, the perceived benefits of physical activity *postpartum* were, in order of importance: weight control, staying fit, and improving mood²¹⁷. The importance that women placed on weight loss postpartum is in contrast to the priority that they placed on management of mood as a motivator for exercise in pregnancy²¹⁷.

Women have reported all of the following barriers to physical activity postpartum: lack of time¹⁶⁰, hours worked outside of the home²⁴¹, finding childcare^{241,266}, presence of other children in the home²⁴¹, lack of opportunity¹⁶⁰, other household responsibilities²⁶¹, and fatigue¹⁶⁰. Again, these responses are different from those in pregnancy when physical limitations are the major barrier.

In terms of additional approaches, a promising intervention was studied among 85 Finnish women who received dietary and physical activity counselling by public health nurses at baby clinics²⁶⁷. Participants kept a food and physical activity record during the

Maternal Weight

eight-month intervention. Although 50% of the intervention group (compared with 30% of controls) returned to their pre-pregnancy weight, the difference did not reach statistical significance.

It is important to note that none of the interventions studied has had a negative impact on breastfeeding²⁶¹.

Motivations and Barriers to Change

In designing an effective intervention it is important to understand women's motivations and barriers to healthy eating and physical activity postpartum. As women adjust to the role of mother, self-care may become less of a priority than caring for children. As such, even when women have adopted or continue to practice healthy eating and physical activity during pregnancy, these practices may not continue postpartum²²⁰.

During the early postpartum period, women may be particularly vulnerable to body-dissatisfaction. In a study of 90 primarily Caucasian women (mean±SD BMI four months postpartum of 25.7±4.8), the proportion of women who reported dieting because they were unhappy with their weight increased from 53% pre-pregnancy to 70% postpartum²²¹. Also, dieting pre-pregnancy was associated with greater body dissatisfaction postpartum²²¹.

In a small study of 22 Black women who were postpartum, the following were identified by them as desirable in a postpartum weight loss intervention: low cost (free), easily accessible geographically, structured, group- or mentor-oriented, including learning good cooking habits, involving the whole family, including child-care, and being staffed by people knowledgeable about the stressors of new motherhood and postpartum depression²⁶⁶.

Screening for Cardiovascular Risk Factors

The 2006 Canadian clinical practice guidelines on the management and prevention of obesity in adults and children recommend for all adults, measurement of BMI and waist circumference¹⁴. For individuals with a BMI>25 kg/m² or a waist circumference above the ethnic specific waist circumference cut-off, the following are also recommended: blood pressure, heart rate, fasting glucose, and fasting lipid profile measurement at regular intervals. Additional investigations, such as liver enzyme tests, urinalysis, and sleep studies are suggested when appropriate.

VIII. Role Modelling to Prevent Childhood Obesity

Children with at least one parent who is obese are at increased risk of becoming obese themselves²⁶⁸. This is intuitive when one considers that parents determine food availability in the home, how the food is eaten in the home, what sort of recreational opportunities are available, children's allowances; and countless other rules and policies that influence the extent to which various members of the family engage in healthful eating and physical activity²⁶⁸.

Maternal Weight

For infants, toddlers, and young children, the mothers' child-feeding practices relate to children's food preferences, energy intake, and ability to sense hunger and fullness^{269,270,271}. Breastfeeding may have a small protective effect for preventing childhood obesity²⁷². Children who are encouraged to eat in response to hunger, and to stop eating when full, are better able to regulate their eating when presented with large amounts of favourite foods in an unsupervised environment^{269,270}. Maternal restriction of high-fat, -salt, or -sugar foods may actually lead to increased BMI and adiposity^{265,266,273} by reinforcing the preference for these 'forbidden foods'^{265,267,274}.

Parents are also important role models for healthy eating^{269,270,274}. Children eat what they see their parents eat and they prefer foods that are offered in positive environments, such as pleasant family meals²⁷⁴. Children who are exposed at an earlier age to vegetables and fruit *enjoy* vegetables and fruit more, and *eat more* vegetables and fruit^{274,275}.

Parents and other care-givers are also important role models for physical activity. The Framingham Children's study found that children of physically active mothers were twice as likely to be active, and those with two active parents were almost six times more likely to be physically active, compared with children of two sedentary parents²⁷⁶.

It follows logically that many groups recommend family-based interventions for prevention (and treatment) of childhood obesity^{14,268,272,277,278,279,280,281,282,283,284}.

In summary, supporting healthy lifestyles amongst postpartum women may support healthy lifestyles for children. Doing so may reinforce for mothers (and fathers) their critical role as role models for healthy eating and lifestyle, and may aid them in supporting their children in developing healthy relationships with food and physical activity.

IX. Conclusions

Obstetric care providers have an important role to play in promoting healthy weight for women in pregnancy and beyond. Women who are overweight or obese pre-pregnancy and those who gain excessive weight during pregnancy, are at increased risk of complications both for themselves and for their fetus/ infant, in a dose-response fashion. The risks associated with maternal weight are comparable to those of other antenatal risk factors, including moderate smoking. Therefore, pregnant women who are overweight or obese should be regarded as higher risk.

There are recognized non-pregnancy related health benefits to having a healthy weight (for example reduced risk of type 2 diabetes). Also, there are barriers present when planning a pregnancy (for example, 50% of pregnancies are unplanned). It is unrealistic to think that this multifaceted health concern will be solved by pre-pregnancy counselling and weight reduction alone. Therefore, it is recommended that steps be taken to achieve a healthy weight outside of pregnancy in addition to targeting women during pregnancy (by promoting weight gain within the IOM guidelines) followed by postpartum interventions to promote the achievement of healthy weights.

Women who are overweight, obese, and/or have had excessive gestational weight gain should be 'flagged' for health care providers who will be assuming or resuming care. These women are at increased risk of complications in the next pregnancy, as well as long-term maternal and paediatric health problems.

Maternal Weight

Given the central role of eating habits in weight loss programs, active involvement of a dietitian should be sought. Consideration could be given to targeting those women who are at greatest risk of postpartum weight retention (i.e. those who are obese, or who gained an excessive amount of weight in pregnancy), and to factoring in their motivations and barriers to healthy lifestyle adoption (i.e. readiness for change).

The findings presented in this paper should serve to motivate the development of better health promotion focussed, women-centred interventions based on published successes and failures with respect to supporting women to experience healthy weights during the childbearing years. Such interventions should also be based on evidence that explores women's motivations and barriers to adopting lifestyles that promote a healthy weight.

Herein is a checklist of better/ promising practices for primary obstetrical healthcare practitioners to support women who are overweight or obese pre-pregnancy, during pregnancy and postpartum, with a goal to improve maternal and perinatal outcomes.

Appendix A: Gestational Weight Gain Recommendations

Table 1: Recommended gestational weight gain relative to pre-pregnancy body mass index (BMI), for singleton pregnancies²⁸⁵

Pre-Pregnancy BMI (kg/m ²)	TOTAL Weight Gain		RATE of Weight Gain
	(kg)	(lb)	
<20	12.5-18	28-40	First trimester gain 1.0 – 3.5 kg (2 – 8 lb). Second and third trimesters gain approximately 0.5kg (1.0lb) per week.
20-27	11.5-16	25-35	First trimester gain 1.0 – 3.5 kg (2 – 8 lb). Second and third trimesters gain approximately 0.4kg (0.75lb) per week.
>27	7-11.5	15-25	First trimester gain 1.0 – 3.5 kg (2 – 8 lb). Second and third trimesters gain approximately 0.3kg (0.5lb) per week.

Appendix B: Pregnancy Care Checklist

Table 2: Pregnancy Care Checklist for Women who are Overweight and Obese

PRE-PREGNANCY

Encourage moderate weight loss prior to conception by physical activity and healthy eating following Eating Well with Canada's Food Guide¹⁸¹ and Canada's Physical Activity Guide to Healthy Active Living¹⁸²

Recommend contraception while aiming for target weight

Counsel about risks associated with maternal overweight and obesity

Optimize glycemic control prior to conception for women with pre-gestational diabetes

Initiate folic acid supplementation 3 months prior to conception among women who are obese

If the woman has undergone bariatric surgery prior to conception, supplementation with vitamin B12, folate, iron and calcium may be indicated

IN PREGNANCY

At initial antenatal visit:

- Record height, weight and BMI (kg/m²)
- Discuss Health Canada weight gain guidelines, set weight gain GOAL and RATE
- Counsel about maternal and fetal risks of overweight/obesity
- Counsel about maternal and fetal risks of excessive gestational weight gain during pregnancy
- Test early glucose tolerance to screen for pre-existing, undiagnosed Type 2 diabetes, especially in the presence of risk factors such as: family history of diabetes, previous GDM, and/or a previous macrosomic infant
- Promote and support breastfeeding

In early pregnancy:

- Confirm dating with early ultrasound
- Conduct detailed anomaly scan at 18 weeks
- Conduct early assessment of liver function, renal function, degree of proteinuria, and ECG to screen for other medical problems
- Promote and support breastfeeding

Throughout or later in pregnancy:

- Measure blood pressure with an appropriately sized cuff
- Monitor weight gain and rate of gain throughout pregnancy, relative to GOAL
- Encourage healthy eating following Eating Well with Canada's Food Guide¹⁸¹
- Encourage moderate physical activity unless there is a (are) contraindication(s)
- Screen for GDM at 24-28 weeks
- Promote and support breastfeeding

Maternal Weight

In preparation for labour and delivery:

- Involve a multidisciplinary team
- Arrange an anaesthetic consult prior to the onset of labour†
- Do complex care planning to ensure appropriate equipment is available (e.g. operating room bed, commodes) and prepare for potential labour complications (including failed induction, slow progress, shoulder dystocia, fetal distress, emergency Caesarean section, thromboembolism postpartum, and post-operative wound infection)
 - Prescribe antibiotic prophylaxis for caesarean section: 2g cephalothin iv if weight is >70kg
 - Administer uterotonics in a timely fashion given increased risk of PPH
- Consider postoperative DVT prophylaxis
- Promote and support breastfeeding

FOLLOWING PREGNANCY

- Counsel regarding healthy eating and physical activity following Eating Well with Canada's Food Guide¹⁸¹ and Canada's Physical Activity Guide to Healthy Active Living¹⁸², promotion of a healthy weight, and other health concerns prior to attempting future pregnancies
- Screen periodically for diabetes and other cardiovascular risk factors according to published guidelines
- Advise on long-term risks of obesity, hypertension and diabetes
- Refer to community support services for the promotion of a healthy lifestyle
- Promote and support breastfeeding

* *Based on expert opinion*^{180,286}

† *A reasonable criterion would be a BMI >35kg/m² but different centres will establish different cut-offs based on logistics.*

GDM (gestational diabetes), DVT (deep vein thrombosis), PPH (postpartum haemorrhage)

Appendix C: References

- ¹ Yeh J, Shelton JA. Increasing prepregnancy body mass index: analysis of trends and contributing variables. *American Journal of Obstetrics & Gynecology*. 2005 Dec;193(6):1994-8.
- ² Ogden CL, Flegal KM, Carroll MD, Johnson CL. Prevalence and trends in overweight among US children and adolescents, 1999-2000. *JAMA*. 2002 Oct 9;288(14):1728-32.
- ³ Okosun et al. Abdominal adiposity in U.S. adults: prevalence and trends, 1960-2000. *Preventive Medicine*. 2004 Jul;39(1):197-206.
- ⁴ Flegal KM, Carroll MD, Ogden CL, Johnson CL. Prevalence and trends in obesity among US adults, 1999-2000. *JAMA*. 2002 Oct 9;288(14):1723-7.
- ⁵ Rocchini AP. Childhood obesity and a diabetes epidemic. *New England Journal of Medicine*. 2002 Mar 14;346(11):854-5.
- ⁶ Kanagalingam MG, Forouhi NG, Greer IA, Sattar N. Changes in booking body mass index over a decade: retrospective analysis from a Glasgow Maternity Hospital. *BJOG*. 2005 Oct;112(10):1431-3.
- ⁷ Sebire NJ et al. Maternal obesity and pregnancy outcome: a study of 287,213 pregnancies in London. *International Journal of Obesity & Related Metabolic Disorders: Journal of the International Association for the Study of Obesity*. 2001 Aug;25(8):1175-82.
- ⁸ Surkan PJ, Hsieh C-C, Johansson ALV, Dickman PW, Cnattingius S. Reasons for increasing trends in large for gestational age births. *Obstetrics & Gynecology*. [Research Support, Non-U.S. Gov't]. 2004 Oct;104(4):720-6.
- ⁹ Callaway LK, Prins JB, Chang AM, McIntyre HD. The prevalence and impact of overweight and obesity in an Australian obstetric population. *Medical Journal of Australia*. 2006 Jan 16;184(2):56-9.
- ¹⁰ Mendez MA, Monteiro CA, Popkin BM. Overweight exceeds underweight among women in most developing countries. *American Journal of Clinical Nutrition*. 2005 Mar;81(3):714-21.
- ¹¹ Katzmarzyk PT. The Canadian obesity epidemic: an historical perspective. *Obesity Research*. 2002 Jul;10(7):666-74.
- ¹² 2004 Canadian Community Health Survey 2.2, Nutrition as reported in Canada's Nutrition and Health Atlas. Accessed August 14, 2008. <http://www.hc-sc.gc.ca/fn-an/surveill/atlas/maps-carte/index-eng.php>
- ¹³ British Columbia Perinatal Database Registry. 2005/6 Pregnancy Health Weights Initiative Report. Prepared December 11, 2007. British Columbia Perinatal Health Program. Unpublished report.

-
- ¹⁴ Lau DC, Douketis JD, Morrison KM, Hramiak IM, Sharma AM, Ur E. Obesity Canada Clinical Practice Guidelines Expert Panel. 2006 Canadian clinical practice guidelines on the management and prevention of obesity in adults and children. *CMAJ*. 2007 Apr 10;176(8):S1-13.
- ¹⁵ Health Canada. Canadian Guidelines for Body Weight Classification in Adults. Ottawa: Minister of Public Works and Government Services Canada; 2003.
- ¹⁶ Janssen I, Katzmarzyk PT, Ross R. Waist circumference and not body mass index explains obesity-related health risk. *American Journal of Clinical Nutrition*. 2004 Mar;79(3):379-84.
- ¹⁷ Janssen I, Katzmarzyk PT, Ross R. Body mass index, waist circumference, and health risk: evidence in support of current National Institutes of Health guidelines. *Archives of Internal Medicine*. 2002 Oct 14;162(18):2074-9.
- ¹⁸ Simpson JA, MacInnis RJ, Peeters A, Hopper JL, Giles GG, English DR. A comparison of adiposity measures as predictors of all-cause mortality: the Melbourne Collaborative Cohort Study. *Obesity*. 2007 Apr;15(4):994-1003.
- ¹⁹ World Health Organization. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet*. 2004 Jan 10;363(9403):157-63).
- ²⁰ Robinson HE, O'Connell CM, Joseph KS, McLeod NL. Maternal outcomes in pregnancies complicated by obesity. *Obstetrics & Gynecology*. 2005 Dec;106(6):1357-64.
- ²¹ Baeten JM, Bukusi EA, Lambe M. Pregnancy complications and outcomes among overweight and obese nulliparous women. *American Journal of Public Health*. 2001 Mar;91(3):436-40.
- ²² Jensen DM et al. Pregnancy outcome and prepregnancy body mass index in 2459 glucose-tolerant Danish women. *American Journal of Obstetrics & Gynecology*. 2003 Jul;189(1):239-44.
- ²³ Watkins ML, Rasmussen SA, Honein MA, Botto LD, Moore CA. Maternal obesity and risk for birth defects. *Pediatrics*. 2003 May;111(5 Part 2):1152-8.
- ²⁴ Ehrenberg HM, Mercer BM, Catalano PM. The influence of obesity and diabetes on the prevalence of macrosomia. *American Journal of Obstetrics & Gynecology*. 2004 Sep;191(3):964-8.
- ²⁵ Doherty DA, Magann EF, Francis J, Morrison JC, Newnham JP. Pre-pregnancy body mass index and pregnancy outcomes. *International Journal of Gynaecology & Obstetrics*. 2006 Dec;95(3):242-7.
- ²⁶ Cnattingius S, Bergstrom R, Lipworth L, Kramer MS. Prepregnancy weight and the risk of adverse pregnancy outcomes. *New England Journal of Medicine*. 1998 Jan 15;338(3):147-52.

- ²⁷ Cedergren MI. Maternal morbid obesity and the risk of adverse pregnancy outcome. *Obstetrics & Gynecology*. 2004 Feb;103(2):219-24.
- ²⁸ Abenhaim H, Kinch R, Morin L, Benjamin A, Usher R. Effect of prepregnancy body mass index categories on obstetrical and neonatal outcomes. *Archives of Gynecology and Obstetrics*. 2007;275(1):39-43.
- ²⁹ Rode L, Nilas L, Wojdemann K, Tabor A. Obesity-related complications in Danish single cephalic term pregnancies. *Obstetrics & Gynecology*. 2005 Mar;105(3):537-42.
- ³⁰ Cnattingius S, Lambe M. Trends in smoking and overweight during pregnancy: prevalence, risks of pregnancy complications, and adverse pregnancy outcomes. *Seminars in Perinatology*. 2002 Aug;26(4):286-95.
- ³¹ Bhattacharya S, Campbell DM, Liston WA, Bhattacharya S. Effect of Body Mass Index on pregnancy outcomes in nulliparous women delivering singleton babies. *BMC Public Health*. 2007;7:168-75.
- ³² Smith GCS, Shah I, Pell JP, Crossley JA, Dobbie R. Maternal obesity in early pregnancy and risk of spontaneous and elective preterm deliveries: a retrospective cohort study. *American Journal of Public Health*. 2007 Jan;97(1):157-62.
- ³³ Nohr EA, Bech BH, Vaeth M, Rasmussen KM, Henriksen TB, Olsen J. Obesity, gestational weight gain and preterm birth: a study within the Danish National Birth Cohort. *Paediatric and Perinatal Epidemiology*. 2007 Jan;21(1):5-14.
- ³⁴ Stotland NE, Washington AE, Caughey AB. Prepregnancy body mass index and the length of gestation at term. *American Journal of Obstetrics & Gynecology*. 2007 Oct;197(4):378.e1-5).
- ³⁵ Oddy WH, Li J, Landsborough L, Kendall GE, Henderson S, Downie J. The association of maternal overweight and obesity with breastfeeding duration. *Journal of Pediatrics*. 2006 Aug;149(2):185-91.
- ³⁶ Raatikainen K, Heiskanen N, Heinonen S. Transition from overweight to obesity worsens pregnancy outcome in a BMI-dependent manner. *Obesity*. 2006 Jan;14(1):165-71.
- ³⁷ Stephansson O, Dickman PW, Johansson A, Cnattingius S. Maternal weight, pregnancy weight gain, and the risk of antepartum stillbirth. *American Journal of Obstetrics & Gynecology*. 2001 Feb;184(3):463-9.
- ³⁸ Kristensen J, Vestergaard M, Wisborg K, Kesmodel U, Secher NJ. Pre-pregnancy weight and the risk of stillbirth and neonatal death. *BJOG*. 2005 Apr;112(4):403-8.
- ³⁹ Waller DK et al. Prepregnancy obesity as a risk factor for structural birth defects. *Archives of Pediatrics & Adolescent Medicine*. 2007 Aug;161(8):745-50.
- ⁴⁰ Dietz PM, Callaghan WM, Morrow B, Cogswell ME. Population-based assessment of the risk of primary cesarean delivery due to excess prepregnancy weight among nulliparous women delivering term infants. *Maternal & Child Health Journal*. 2005 Sep;9(3):237-44.

- ⁴¹ Dempsey JC, Ashiny Z, Qiu C-F, Miller RS, Sorensen TK, Williams MA. Maternal pre-pregnancy overweight status and obesity as risk factors for cesarean delivery. *Journal of Maternal-Fetal & Neonatal Medicine*. 2005 Mar;17(3):179-85.
- ⁴² Barau G et al. Linear association between maternal pre-pregnancy body mass index and risk of caesarean section in term deliveries. *BJOG*. 2006 Oct;113(10):1173-7.
- ⁴³ Sherrard A, Platt RW, Vallerand D, Usher RH, Zhang X, Kramer MS. Maternal anthropometric risk factors for caesarean delivery before or after onset of labour. *BJOG: An International Journal of Obstetrics & Gynaecology*. 2007 Sep;114(9):1088-96.
- ⁴⁴ Cedergren M, Kallen B. Maternal obesity and the risk for orofacial clefts in the offspring. *Cleft Palate-Craniofacial Journal*. 2005 Jul;42(4):367-71.
- ⁴⁵ Rich-Edwards JW et al. Adolescent body mass index and infertility caused by ovulatory disorder. *American Journal of Obstetrics & Gynecology*. 1994 Jul;171(1):171-7.
- ⁴⁶ Pasquali R, Gambineri A, Pagotto U. The impact of obesity on reproduction in women with polycystic ovary syndrome. *BJOG*. 2006 Oct;113(10):1148-59.
- ⁴⁷ Galtier-Dereure F, Pujol P, Dewailly D, Bringer J. Choice of stimulation in polycystic ovarian syndrome: the influence of obesity. *Human Reproduction*. 1997 Oct;12 Suppl 1:88-96.
- ⁴⁸ Bellver J, Busso C, Pellicer A, Remohi J, Simon C. Obesity and assisted reproductive technology outcomes. *Reproductive Biomedicine Online*. 2006 May;12(5):562-8.
- ⁴⁹ Fedorcsak P et al. Impact of overweight and underweight on assisted reproduction treatment. *Human Reproduction*. 2004 Nov;19(11):2523-8.
- ⁵⁰ van Swieten ECAM, van der Leeuw-Harmsen L, Badings EA, van der Linden PJQ. Obesity and Clomiphene Challenge Test as predictors of outcome of in vitro fertilization and intracytoplasmic sperm injection. *Gynecologic & Obstetric Investigation*. 2005;59(4):220-4.
- ⁵¹ Maheshwari A, Stofberg L, Bhattacharya S. Effect of overweight and obesity on assisted reproductive technology-a systematic review. *Human Reproduction Update*. 2007 Sep-Oct;13(5):433-44.
- ⁵² Rosenberg L, Palmer JR, Wise LA, Horton NJ, Kumanyika SK, Adams-Campbell LL. A prospective study of the effect of childbearing on weight gain in African-American women. *Obesity Research*. 2003 Dec;11(12):1526-35.
- ⁵³ Chu SY et al. Maternal obesity and risk of gestational diabetes mellitus. *Diabetes Care*. 2007 Aug;30(8):2070-6.
- ⁵⁴ Jovanovic L. What is so bad about a big baby?[see comment][comment]. *Diabetes Care*. 2001 Aug;24(8):1317-8.

- ⁵⁵ Goldman M, Kitzmiller JL, Abrams B, Cowan RM, Laros Jr RK. Obstetric complications with GDM: effects of maternal weight. *Diabetes*. 1991 Dec;40 Suppl 2:79-82.
- ⁵⁶ Yogev Y, Xenakis EMJ, Langer O. The association between preeclampsia and the severity of gestational diabetes: the impact of glycemic control. *American Journal of Obstetrics & Gynecology*. 2004 Nov;191(5):1655-60.
- ⁵⁷ Catalano PM, Kirwan JP, Haugel-de Mouzon S, King J. Gestational diabetes and insulin resistance: role in short- and long-term implications for mother and fetus. *Journal of Nutrition*. 2003 May;133(5 Suppl 2):1674S-83S.
- ⁵⁸ Lauenborg J et al. Increasing incidence of diabetes after gestational diabetes: a long-term follow-up in a Danish population. *Diabetes Care*. 2004 May;27(5):1194-9.
- ⁵⁹ Lauenborg J et al. The prevalence of the metabolic syndrome in a danish population of women with previous gestational diabetes mellitus is three-fold higher than in the general population. *Journal of Clinical Endocrinology & Metabolism*. 2005 Jul;90(7):4004-10.
- ⁶⁰ Kjos SL, Peters RK, Xiang A, Henry OA, Montoro M, Buchanan TA. Predicting future diabetes in Latino women with gestational diabetes. Utility of early postpartum glucose tolerance testing. *Diabetes*. 1995 May;44(5):586-91.
- ⁶¹ Rosenberg TJ, Garbers S, Chavkin W, Chiasson MA. Prepregnancy weight and adverse perinatal outcomes in an ethnically diverse population. *Obstetrics & Gynecology*. 2003 Nov;102(5 Pt 1):1022-7.
- ⁶² Weiss JL et al. Obesity, obstetric complications and Caesarean delivery rate: a population-based screening study. *American Journal of Obstetrics & Gynecology*. 2004 Apr;190(4):1091-7.
- ⁶³ O'Brien TE, Ray JG, Chan W-S. Maternal body mass index and the risk of preeclampsia: a systematic overview. *Epidemiology*. 2003 May;14(3):368-74.
- ⁶⁴ Edwards LE, Hellerstedt WL, Alton IR, Story M, Himes JH. Pregnancy complications and birth outcomes in obese and normal-weight women: effects of gestational weight change. *Obstetrics & Gynecology*. 1996 Mar;87(3):389-94.
- ⁶⁵ Andreasen KR, Andersen ML, Schantz AL. Obesity and pregnancy.[see comment]. *Acta Obstetrica et Gynecologica Scandinavica*. 2004 Nov;83(11):1022-9.
- ⁶⁶ Larsen TB, Sorensen HT, Gislum M, Johnsen SP. Maternal smoking, obesity, and risk of venous thromboembolism during pregnancy and the puerperium: a population-based nested case-control study. *Thrombosis Research*. 2007;120(4):505-9.
- ⁶⁷ Lewis G. The Confidential Enquiry into Maternal and Child Health (CEMACH). *Saving Mothers' Lives: reviewing maternal deaths to make motherhood safer - 2003-2005. The Seventh Report on Confidential Enquiries into Maternal Deaths in the United Kingdom*. London. 2007.

- ⁶⁸ Kabiru W, Raynor BD. Obstetric outcomes associated with increase in BMI category during pregnancy. *American Journal of Obstetrics & Gynecology*. 2004 Sep;191(3):928-32.
- ⁶⁹ Sheiner E, Levy A, Menes TS, Silverberg D, Katz M, Mazor M. Maternal obesity as an independent risk factor for caesarean delivery. *Paediatric and Perinatal Epidemiology*. 2004 May;18(3):196-201.
- ⁷⁰ Crane SS, Wojtowycz MA, Dye TD, Aubry RH, Artal R. Association between pre-pregnancy obesity and the risk of cesarean delivery. *Obstetrics & Gynecology*. 1997 Feb;89(2):213-6.
- ⁷¹ Perlow JH, Morgan MA, Montgomery D, Towers CV, Porto M. Perinatal outcome in pregnancy complicated by massive obesity. *American Journal of Obstetrics & Gynecology*. 1992 Oct;167(4 Pt 1):958-62.
- ⁷² Garbacia Jr JA, Richter M, Miller S, Barton JJ. Maternal weight and pregnancy complications. *American Journal of Obstetrics & Gynecology*. 1985 May 15;152(2):238-45.
- ⁷³ Michlin R et al. Maternal obesity and pregnancy outcome. *Israel Medical Association Journal*. 2000 Jan;2(1):10-3.
- ⁷⁴ Durnwald CP, Ehrenberg HM, Mercer BM. The impact of maternal obesity and weight gain on vaginal birth after cesarean section success. *American Journal of Obstetrics & Gynecology*. 2004 Sep;191(3):954-7.
- ⁷⁵ Chu SY et al. Maternal obesity and risk of cesarean delivery: a meta-analysis. *Obesity Reviews*. 2007 Sept; 8(5): 385-94.
- ⁷⁶ Kaiser PS, Kirby RS. Obesity as a risk factor for cesarean in a low-risk population. *Obstetrics & Gynecology*. 2001 Jan;97(1):39-43.
- ⁷⁷ Beattie PG, Rings TR, Hunter MF, Lake Y. Risk factors for wound infection following caesarean section. *Australian & New Zealand Journal of Obstetrics & Gynaecology*. 1994 Aug;34(4):398-402.
- ⁷⁸ Martens MG, Kolrud BL, Faro S, Maccato M, Hammill H. Development of wound infection or separation after cesarean delivery: prospective evaluation of 2,431 cases. *Journal of Reproductive Medicine*. 1995 Mar;40(3):171-5.
- ⁷⁹ Formiguera X, Canton A. Obesity: epidemiology and clinical aspects. *Best Practice & Research in Clinical Gastroenterology*. 2004 Dec;18(6):1125-46.
- ⁸⁰ Bianco AT, Smilen SW, Davis Y, Lopez S, Lapinski R, Lockwood CJ. Pregnancy outcome and weight gain recommendations for the morbidly obese woman. *Obstetrics & Gynecology*. 1998 Jan;91(1):97-102.
- ⁸¹ Bodnar LM, Siega-Riz AM, Cogswell ME. High prepregnancy BMI increases the risk of postpartum anemia. *Obesity Research*. [Research Support, U.S. Gov't, P.H.S.]. 2004 Jun;12(6):941-8.

- ⁸² Hilson JA, Rasmussen KM, Kjolhede CL. Excessive weight gain during pregnancy is associated with earlier termination of breast-feeding among White women. *Journal of Nutrition*. 2006 Jan;136(1):140-6.
- ⁸³ Donath SM, Amir LH. Does maternal obesity adversely affect breastfeeding initiation and duration? *Breastfeeding Review*. 2000 Nov;8(3):29-33.
- ⁸⁴ Hilson JA, Rasmussen KM, Kjolhede CL. Maternal obesity and breast-feeding success in a rural population of white women. *American Journal of Clinical Nutrition*. 1997 Dec;66(6):1371-8.
- ⁸⁵ Li R, Jewell S, Grummer-Strawn L. Maternal obesity and breast-feeding practices. *American Journal of Clinical Nutrition*. 2003 Apr;77(4):931-6.
- ⁸⁶ Amir LH, Donath S. A systematic review of maternal obesity and breastfeeding intention, initiation and duration. *BMC Pregnancy & Childbirth*. 2007;7:9-22.
- ⁸⁷ Wang JX, Davies MJ, Norman RJ. Polycystic ovarian syndrome and the risk of spontaneous abortion following assisted reproductive technology treatment. *Human Reproduction*. 2001 Dec;16(12):2606-9.
- ⁸⁸ Lashen H, Fear K, Sturdee DW. Obesity is associated with increased risk of first trimester and recurrent miscarriage: matched case-control study. *Human Reproduction*. 2004 Jul;19(7):1644-6.
- ⁸⁹ Bellver J et al. Obesity and the risk of spontaneous abortion after oocyte donation. *Fertility & Sterility*. 2003 May;79(5):1136-40.
- ⁹⁰ Chu SY et al. Maternal obesity and risk of stillbirth: a metaanalysis. *American Journal of Obstetrics & Gynecology*. 2007 Sep;197(3):223-8.
- ⁹¹ Nohr EA, Bech BH, Davies MJ, Frydenberg M, Henriksen TB, Olsen J. Prepregnancy obesity and fetal death: a study within the Danish National Birth Cohort. *Obstetrics & Gynecology*. 2005 Aug;106(2):250-9.
- ⁹² Waller DK et al. Are obese women at higher risk for producing malformed offspring? *American Journal of Obstetrics & Gynecology*. 1994 Feb;170(2):541-8.
- ⁹³ Werler MM, Louik C, Shapiro S, Mitchell AA. Prepregnant weight in relation to risk of neural tube defects. *JAMA*. 1996 Apr 10;275(14):1089-92.
- ⁹⁴ Watkins ML, Scanlon KS, Mulinare J, Khoury MJ. Is maternal obesity a risk factor for anencephaly and spina bifida? *Epidemiology*. 1996 Sep;7(5):507-12.
- ⁹⁵ Shaw GM, Todoroff K, Schaffer DM, Selvin S. Maternal height and prepregnancy body mass index as risk factors for selected congenital anomalies. *Paediatric and Perinatal Epidemiology*. 2000 Jul;14(3):234-9.
- ⁹⁶ Shaw GM, Velie EM, Schaffer D. Risk of neural tube defect-affected pregnancies among obese women. *JAMA*. 1996 Apr 10;275(14):1093-6.

- ⁹⁷ Anderson JL, Waller DK, Canfield MA, Shaw GM, Watkins ML, Werler MM. Maternal obesity, gestational diabetes, and central nervous system birth defects. *Epidemiology*. 2005 Jan;16(1):87-92.
- ⁹⁸ Moore LL, Singer MR, Bradlee ML, Rothman KJ, Milunsky A. A prospective study of the risk of congenital defects associated with maternal obesity and diabetes mellitus. *Epidemiology*. 2000 Nov;11(6):689-94.
- ⁹⁹ Shaw GM, Todoroff K, Finnell RH, Lammer EJ. Spina bifida phenotypes in infants or fetuses of obese mothers. *Teratology*. 2000 May;61(5):376-81.
- ¹⁰⁰ Hendricks KA, Nuno OM, Suarez L, Larsen R. Effects of hyperinsulinemia and obesity on risk of neural tube defects among Mexican Americans. *Epidemiology*. 2001 Nov;12(6):630-5.
- ¹⁰¹ Dandolu V, Jain NJ, Hernandez E, Kruse L. Shoulder dystocia at noninstrumental vaginal delivery. *American Journal of Perinatology*. 2006 Oct;23(7):439-44.
- ¹⁰² Hediger ML, Overpeck MD, Maurer KR, Kuczmarski RJ, McGlynn A, Davis WW. Growth infants and young children born small or large for gestational age: findings from the Third National Health and Nutrition Examination Survey. *Archives of Pediatrics & Adolescent Medicine*. 1998 Dec; 152(12):1225-31.
- ¹⁰³ Kallen K. The impact of maternal smoking during pregnancy on delivery outcome. *European Journal of Public Health*. 2001 Sep;11(3):329-33.
- ¹⁰⁴ Kesmodel U, Wisborg K, Olsen SF, Henriksen TB, Secher NJ. Moderate alcohol intake during pregnancy and the risk of stillbirth and death in the first year of life. *American Journal of Epidemiology*. 2002 Feb 15;155(4):305-12.
- ¹⁰⁵ Hernandez-Diaz S, Werler MM, Walker AM, Mitchell AA. Neural tube defects in relation to use of folic acid antagonists during pregnancy. *American Journal of Epidemiology*. 2001 May 15;153(10):961-8.
- ¹⁰⁶ O'Brien E et al. Practice guidelines of the European Society of Hypertension for clinic, ambulatory and self blood pressure measurement. *Journal of Hypertension*. 2005 Apr;23(4):697-701.
- ¹⁰⁷ Wolfe HM, Sokol RJ, Martier SM, Zador IE. Maternal obesity: a potential source of error in sonographic prenatal diagnosis. *Obstetrics & Gynecology*. 1990 Sep;76(3 Pt 1):339-42.
- ¹⁰⁸ Hendler I, Blackwell SC, Treadwell MC, Bujold E, Sokol RJ, Sorokin Y. Does advanced ultrasound equipment improve the adequacy of ultrasound visualization of fetal cardiac structures in the obese gravid woman? *American Journal of Obstetrics & Gynecology*. 2004 Jun;190(6):1616-9; discussion 9-20.
- ¹⁰⁹ Krishnamoorthy U, Schram CMH, Hill SR. Maternal obesity in pregnancy: Is it time for meaningful research to inform preventive and management strategies? *BJOG*. 2006 Oct;113(10):1134-40.

- ¹¹⁰ Joint WHO/FAO Expert Consultation. Diet, Nutrition and the Prevention of Chronic Diseases. 2003. Accessed Jan 27, 2009. http://whqlibdoc.who.int/trs/WHO_TRS_916.pdf.
- ¹¹¹ Jain MG et al. Body mass index and mortality in women: follow-up of the Canadian National Breast Screening Study cohort. *International Journal of Obesity (London)*. 2005 Jul;29(7):792-7.
- ¹¹² McTigue K, et al. Mortality and cardiac and vascular outcomes in extremely obese women. *JAMA*. 2006 Jul 5;296(1):79-86.
- ¹¹³ Koster A et al. The combined relations of adiposity and smoking on mortality. *American Journal of Clinical Nutrition*. 2008;88:1206-1212.
- ¹¹⁴ Jensen MK, Chiuve SE, Rimm EB, Dethlefsen C, Tjønneland A, Joensen AM, Overvad K. Obesity, behavioral lifestyle factors, and risk of acute coronary events. *Circulation*. 2008 Jun 17;117(24):3062-9.
- ¹¹⁵ Wang Y et al. Association between obesity and kidney disease: a systematic review and meta-analysis. *Kidney International*. 2008 Jan;73(1):19-33.
- ¹¹⁶ Elsayed EF et al. Waist-to-hip ratio, body mass index, and subsequent kidney disease and death. *American Journal of Kidney Disease*. 2008 Jul;52(1):29-38.
- ¹¹⁷ Renehan AG et al. Body-mass index and incidence of cancer: a systematic review and meta-analysis of prospective observational studies. *Lancet*. 2008 Feb 16;371(9612):569-78.
- ¹¹⁸ Beuther DA et al. Overweight, obesity, and incident asthma: a meta-analysis of prospective epidemiologic studies. *American Journal of Respiratory Critical Care Medicine*. 2007 Apr 1;175(7):661-6.
- ¹¹⁹ Messier SP et al. Exercise and dietary weight loss in overweight and obese older adults with knee osteoarthritis: the arthritis, diet, and activity promotion trial. *Arthritis & Rheumatism*. 2004 May;50(5):1501-1510.
- ¹²⁰ Flugsrud GB, Nordsletten L, Espehaug B, Havelin LI, Engeland A, Meyer HE. The impact of body mass index on later total hip arthroplasty for primary osteoarthritis: a cohort study in 1.2 million persons. *Arthritis & Rheumatism*. 2006 Mar;54(3):802-807.
- ¹²¹ Gorospe EC, Dave JK. The risk of dementia with increased body mass index. *Age & Ageing*. 2007;36:23-29.
- ¹²² Salsberry PJ, Reagan PB. Dynamics of early childhood overweight. *Pediatrics*. 2005 Dec;116(6):1329-38.
- ¹²³ Laitinen J, Power C, Jarvelin MR. Family social class, maternal body mass index, childhood body mass index, and age at menarche as predictors of adult obesity. *American Journal of Clinical Nutrition*. 2001 Sep;74(3):287-94.
- ¹²⁴ Hunter WA et al. Insulin sensitivity in the offspring of women with type 1 and type 2 diabetes. *Diabetes Care*. 2004 May;27(5):1148-52.

- ¹²⁵ Boney CM, Verma A, Tucker R, Vohr BR. Metabolic syndrome in childhood: association with birth weight, maternal obesity, and gestational diabetes mellitus. *Pediatrics*. 2005 Mar;115(3):e290-6.
- ¹²⁶ Gillman MW, Rifas-Shiman S, Berkey CS, Field AE, Colditz GA. Maternal gestational diabetes, birth weight, and adolescent obesity. *Pediatrics*. 2003 Mar;111(3):e221-6.
- ¹²⁷ Pettitt DJ, Knowler WC. Long-term effects of the intrauterine environment, birth weight, and breast-feeding in Pima Indians. *Diabetes Care*. 1998 Aug;21 Suppl 2:B138-41.
- ¹²⁸ Silverman BL, Rizzo TA, Cho NH, Metzger BE. Long-term effects of the intrauterine environment. The Northwestern University Diabetes in Pregnancy Center. *Diabetes Care*. 1998 Aug;21 Suppl 2:B142-9.
- ¹²⁹ Weintrob N, Karp M, Hod M. Short- and long-range complications in offspring of diabetic mothers. *Journal of Diabetes & its Complications*. 1996 Sep-Oct;10(5):294-301.
- ¹³⁰ Manderson JG, Mullan B, Patterson CC, Hadden DR, Traub AI, McCance DR. Cardiovascular and metabolic abnormalities in the offspring of diabetic pregnancy. *Diabetologia*. 2002 Jul;45(7):991-6.
- ¹³¹ Touger L, Looker HC, Krakoff J, Lindsay RS, Cook V, Knowler WC. Early growth in offspring of diabetic mothers. *Diabetes Care*. 2005 Mar;28(3):585-9.
- ¹³² Whitaker RC. Predicting preschooler obesity at birth: the role of maternal obesity in early pregnancy. *Pediatrics*. 2004 Jul;114(1):e29-36.
- ¹³³ Levin BE, Govek E. Gestational obesity accentuates obesity in obesity-prone progeny. *American Journal of Physiology*. 1998 Oct;275(4 Pt 2):R1374-9.
- ¹³⁴ Vickers MH, Breier BH, Cutfield WS, Hofman PL, Gluckman PD. Fetal origins of hyperphagia, obesity, and hypertension and postnatal amplification by hypercaloric nutrition. *American Journal of Physiology - Endocrinology & Metabolism*. 2000 Jul;279(1):E83-7.
- ¹³⁵ Hillier TA, Pedula KL, Schmidt MM, Mullen JA, Charles M-A, Pettitt DJ. Childhood obesity and metabolic imprinting: the ongoing effects of maternal hyperglycemia. *Diabetes Care*. 2007 Sep;30(9):2287-92.
- ¹³⁶ Galtier-Dereure F, Boegner C, Bringer J. Obesity and pregnancy: complications and cost. *American Journal of Clinical Nutrition*. 2000 May;71(5 Suppl):1242S-8S.
- ¹³⁷ Galtier-Dereure F, Montpeyroux F, Boulot P, Bringer J, Jaffiol C. Weight excess before pregnancy: complications and cost. *International Journal of Obesity & Related Metabolic Disorders*. 1995 Jul;19(7):443-8.
- ¹³⁸ Chu SY et al. Association between obesity during pregnancy and increased use of health care. *New England Journal of Medicine*. 2008 Apr 3;358(14):1444-53.

-
- ¹³⁹ Institute of Medicine: Subcommittee on Dietary Intake and Nutrient Supplements during Pregnancy, Committee on Nutritional Status during Pregnancy and Lactation, Food and Nutrition Board. Nutrition during pregnancy, weight gain and nutrient supplements: Report of the Subcommittee on Nutritional Status and Weight Gain during Pregnancy. Washington, DC: National Academy Press; 1990:1–233, 333–41.
- ¹⁴⁰ Nutrition during pregnancy. ACOG Technical Bulletin Number 179--April 1993. *International Journal of Gynaecology & Obstetrics*. 43(1):67-74, 1993 Oct.
- ¹⁴¹ Abrams B, Altman SL, Pickett KE. Pregnancy weight gain: still controversial. *American Journal of Clinical Nutrition*. 2000 May;71(5 Suppl):1233S-41S.
- ¹⁴² Thorsdottir I, Torfadottir JE, Birgisdottir BE, Geirsson RT. Weight gain in women of normal weight before pregnancy: complications in pregnancy or delivery and birth outcome. *Obstetrics & Gynecology*. 2002 May;99(5 Pt 1):799-806.
- ¹⁴³ DeVader SR, Neeley HL, Myles TD, Leet TL. Evaluation of gestational weight gain guidelines for women with normal prepregnancy body mass index. *Obstetrics & Gynecology*. 2007 Oct;110(4):745-51.
- ¹⁴⁴ Cedergren MI. Optimal gestational weight gain for body mass index categories.[see comment]. *Obstetrics & Gynecology*. 2007 Oct;110(4):759-64.
- ¹⁴⁵ Schieve LA, Cogswell ME, Scanlon KS. An empiric evaluation of the Institute of Medicine's pregnancy weight gain guidelines by race. *Obstetrics & Gynecology*. 1998 Jun;91(6):878-84.
- ¹⁴⁶ Johnson JW, Longmate JA, Frentzen B. Excessive maternal weight and pregnancy outcome. *American Journal of Obstetrics & Gynecology*. 1992 Aug;167(2):353-70; discussion 70-2.
- ¹⁴⁷ Cedergren M. Effects of gestational weight gain and body mass index on obstetric outcome in Sweden. *International Journal of Gynaecology & Obstetrics*. 2006 Jun;93(3):269-74.
- ¹⁴⁸ Stotland NE, Hopkins LM, Caughey AB. Gestational weight gain, macrosomia, and risk of cesarean birth in nondiabetic nulliparas. *Obstetrics & Gynecology*. 2004 Oct;104(4):671-7.
- ¹⁴⁹ Shepard MJ, Hellenbrand KG, Bracken MB. Proportional weight gain and complications of pregnancy, labor, and delivery in healthy women of normal prepregnant stature. *American Journal of Obstetrics & Gynecology*. 1986 Nov;155(5):947-54.
- ¹⁵⁰ Stotland NE, Cheng YW, Hopkins LM, Caughey AB. Gestational weight gain and adverse neonatal outcome among term infants. *Obstetrics & Gynecology*. 2006 Sep;108(3 Pt 1):635-43.
- ¹⁵¹ Rhodes JC, Schoendorf KC, Parker JD. Contribution of excess weight gain during pregnancy and macrosomia to the cesarean delivery rate, 1990-2000. *Pediatrics*. 2003 May;111(5 Part 2):1181-5.

- ¹⁵² Oken E, Taveras EM, Kleinman KP, Rich-Edwards JW, Gillman MW. Gestational weight gain and child adiposity at age 3 years. *American Journal of Obstetrics & Gynecology*. 2007 Apr;196(4):322.e1-8.
- ¹⁵³ Carmichael S, Abrams B, Selvin S. The pattern of maternal weight gain in women with good pregnancy outcomes. *American Journal of Public Health*. 1997 Dec;87(12):1984-8.
- ¹⁵⁴ Muscati SK, Gray-Donald K, Koski KG. Timing of weight gain during pregnancy: promoting fetal growth and minimizing maternal weight retention. *International Journal of Obesity & Related Metabolic Disorders*. 1996 Jun;20(6):526-32.
- ¹⁵⁵ Siega-Riz AM, Adair LS, Hobel CJ. Institute of Medicine maternal weight gain recommendations and pregnancy outcome in a predominantly Hispanic population. *Obstetrics & Gynecology*. 1994 Oct;84(4):565-73.
- ¹⁵⁶ Dawes MG, Grudzinskas JG. Patterns of maternal weight gain in pregnancy. *British Journal of Obstetrics & Gynaecology*. 1991 Feb;98(2):195-201.
- ¹⁵⁷ Abrams B, Parker JD. Maternal weight gain in women with good pregnancy outcome. *Obstetrics & Gynecology*. 1990 Jul;76(1):1-7.
- ¹⁵⁸ Kinnunen TI, Luoto R, Gissler M, Hemminki E. Pregnancy weight gain from 1960s to 2000 in Finland. *International Journal of Obesity & Related Metabolic Disorders*. 2003 Dec;27(12):1572-7.
- ¹⁵⁹ Gunderson EP, Abrams B. Epidemiology of gestational weight gain and body weight changes after pregnancy. *Epidemiologic Reviews*. 2000;22(2):261-74.
- ¹⁶⁰ Harris HE, Ellison GT, Clement S. Do the psychosocial and behavioral changes that accompany motherhood influence the impact of pregnancy on long-term weight gain? *Journal of Psychosomatic Obstetrics & Gynecology*. 1999 Jun;20(2):65-79.
- ¹⁶¹ National Research Council and Institute of Medicine. *Influence of Pregnancy Weight on Maternal and Child Health: Workshop Report*. Washington, DC: The National Academies Press; 2007
- ¹⁶² Olafsdottir AS, Skuladottir GV, Thorsdottir I, Hauksson A, Steingrimsdottir L. Maternal diet in early and late pregnancy in relation to weight gain. *International Journal of Obesity*. 2006;30:492-499.
- ¹⁶³ Clapp III J. Maternal carbohydrate intake and pregnancy outcome. *Proceedings of the Nutrition Society*. 2002;61(1):45-50.
- ¹⁶⁴ Siega-Riz AM, Hobel CJ. Predictors of poor maternal weight gain from baseline anthropometric, psychosocial, and demographic information in a Hispanic population. *Journal of the American Dietetic Association*. 1997;97:1264-1268.
- ¹⁶⁵ Conway R, Reddy S, Davies J. Dietary restraint and weight gain during pregnancy. *European Journal of Clinical Nutrition*. 1999;53:849-853.

- ¹⁶⁶ Siega-Riz AM, Evenson KR, Doyle N. Pregnancy-related weight gain – a link to obesity? *Nutrition Reviews*. 2004;62(7):S105-S111.
- ¹⁶⁷ World Health Organization. *The Ottawa Charter for Health Promotion*. First International Conference on Health Promotion, Ottawa:1986 Nov. Accessed February 15, 2009. <http://www.who.int/healthpromotion/conferences/previous/ottawa/en/>.
- ¹⁶⁸ British Columbia Ministry of Health Services. *Supporting Local Collaborative Models for Sustainable Maternity Care in British Columbia: Recommendations from the Maternity Care Enhancement Project*. 2004 Dec. Victoria, BC: British Columbia Ministry of Health Services. Accessed February 14, 2009. http://www.health.gov.bc.ca/library/publications/year/2004/mcep_recommend_dec2004.pdf.
- ¹⁶⁹ Wilson GT, Schlam TR. The transtheoretical model and motivational interviewing in the treatment of eating and weight disorders. *Clinical Psychology Review*. 2004;24:361-378.
- ¹⁷⁰ Smith West D, DiLillo V, Bursac Z, Gore SA, Greene PG. Motivational interviewing improves weight loss in women with type 2 diabetes. *Diabetes Care*. 2007;30:1081-1087.
- ¹⁷¹ Schwartz RP et al. Office-based motivational interviewing to prevent childhood obesity: a feasibility study. *Arch Pediatr Adolesc Med*. 2007;161:495-501.
- ¹⁷² Carels RA et al. Using motivational interviewing as a supplement to obesity treatment: a stepped-care approach. *Health Psychology*. 2007;26(3):369-374.
- ¹⁷³ Webber KH, Tate DF, Quintiliani LM. Motivational interviewing in internet groups: a pilot study for weight loss. *Journal of the American Dietetic Association*. 2008;108:1029-1032.
- ¹⁷⁴ Burke BL, Arkowitz H, Menchola M. The efficacy of motivational interviewing: a meta-analysis of controlled clinical trials. *Journal of Consulting and Clinical Psychology*. 2003;71(5):843-861.
- ¹⁷⁵ Resnicow K, Davis R, Rollnick S. Motivational interviewing for pediatric obesity: conceptual issues and evidence review. *Journal of the American Dietetic Association*. 2006;106:2024-2033.
- ¹⁷⁶ Rubak S, Sandbaek A, Lauritzen T, Christensen B. Motivational interviewing: a systematic review and meta-analysis. *British Journal of General Practice*. 2005;55:305-312.
- ¹⁷⁷ Rollnick S, Miller WR, Butler CC. *Motivational Interviewing in Health Care: Helping Patients Change Behavior*. New York: The Guilford Press; 2008.
- ¹⁷⁸ Prochaska JO, Diclemente CC. Stages of change in the modification of problem behaviors. *Prog Behav Modif*. 1992;28:183-218.
- ¹⁷⁹ Yu C, Teoh T, Robinson S. Obesity in pregnancy. *BJOG*. 2006;113:1117–1125

- ¹⁸⁰ Mighty HE, Fahey AJ. Obesity and pregnancy complications. *Current Diabetes Reports*. 2007 Aug;7(4):289-94.
- ¹⁸¹ Health Canada. *Eating Well with Canada's Food Guide: A Resource for Educators and Communicators*. Ottawa: Minister of Health; 2007.
- ¹⁸² Public Health Agency of Canada & Canadian Society for Exercise Physiology. *Handbook for Canada's Physical Activity Guide to Healthy Active Living*. Ottawa; 1998.
- ¹⁸³ BC Perinatal Health Program. *BCRCP Guideline: Tobacco Use in the Perinatal Period*. 2006. Accessed August 14, 2008 <http://www.bcphp.ca/sites/bcrp/files/Guidelines/SubstanceUse/TobaccoGuidelines.pdf>
- ¹⁸⁴ BC Perinatal Health Program. *BCRCP Guidelines for Alcohol Use in the Perinatal Period and Fetal Alcohol Spectrum Disorder*. 2005. Accessed August 14, 2008 <http://www.bcphp.ca/sites/bcrp/files/Guidelines/SubstanceUse/Alcoholguideline.pdf>
- ¹⁸⁵ American College of Obstetricians and Gynecologists. Obesity in pregnancy, opinion no. 315. *Obstet Gynecol*.
- ¹⁸⁶ Pasquali R et al. Clinical and hormonal characteristics of obese amenorrheic hyperandrogenic women before and after weight loss. *Journal of Clinical Endocrinology & Metabolism*. 1989 Jan;68(1):173-9.
- ¹⁸⁷ Kiddy DS et al. Improvement in endocrine and ovarian function during dietary treatment of obese women with polycystic ovary syndrome. *Clinical Endocrinology*. 1992 Jan;36(1):105-11.
- ¹⁸⁸ Crosignani PG, Colombo M, Vegetti W, Somigliana E, Gessati A, Ragni G. Overweight and obese anovulatory patients with polycystic ovaries: parallel improvements in anthropometric indices, ovarian physiology and fertility rate induced by diet. *Human Reproduction*. 2003 Sep;18(9):1928-32.
- ¹⁸⁹ Clark AM et al. Weight loss results in significant improvement in pregnancy and ovulation rates in anovulatory obese women. *Human Reproduction*. 1995 Oct;10(10):2705-12.
- ¹⁹⁰ Patel JA, Colella JJ, Esaka E, Patel NA, Thomas RL. Improvement in infertility and pregnancy outcomes after weight loss surgery. *Medical Clinics of North America*. 2007 May;91(3):515-28.
- ¹⁹¹ Villamor E, Cnattingius S. Interpregnancy weight change and risk of adverse pregnancy outcomes: a population-based study. *Lancet*. 2006 Sep 30;368(9542):1164-70.
- ¹⁹² Woodard CB. Pregnancy following bariatric surgery. *Journal of Perinatal & Neonatal Nursing*. 2004 Oct-Dec;18(4):329-40.
- ¹⁹³ Malinowski SS. Nutritional and metabolic complications of bariatric surgery. *American Journal of the Medical Sciences*. [Review]. 2006 Apr;331(4):219-25.
- ¹⁹⁴ Alvarez-Leite JI. Nutrient deficiencies secondary to bariatric surgery. *Current Opinion in Clinical Nutrition & Metabolic Care*. 2004 Sep;7(5):569-75.

¹⁹⁵ Wilson RD et al. Preconceptional vitamin / folic acid supplementation 2007: the use of folic acid in combination with a multivitamin supplement for the prevention of neural tube defects and other congenital anomalies. *Journal of Obstetrics & Gynaecology Canada*. 2007 Dec; 29(12):1003-26.

¹⁹⁶ Abir F, Bell R. Assessment and management of the obese patient. *Critical Care Medicine*. 2004 Apr;32(4Suppl):S87-91.

¹⁹⁷ Catalano PM. Management of obesity in pregnancy. *Obstetrics & Gynecology*. 2007 Feb;109(2 Pt 1):419-33.

¹⁹⁸ British Columbia Perinatal Health Program. Obstetric Guideline 10A: Gestational Diabetes. 2001 Oct. Accessed February 3, 2009. <http://www.bcphp.ca/sites/bcrp/files/Guidelines/Obstetrics/MasterGestationalDiabetesOctober.pdf>.

¹⁹⁹ Cogswell ME, Scanlon KS, Fein SB, Schieve LA. Medically advised, mother's personal target, and actual weight gain during pregnancy. *Obstetrics & Gynecology*. 1999 Oct;94(4):616-22.

²⁰⁰ Stotland NE, Haas JS, Brawarsky P, Jackson RA, Fuentes-Afflick E, Escobar GJ. Body mass index, provider advice, and target gestational weight gain. *Obstetrics & Gynecology*. 2005 Mar;105(3):633-8.

²⁰¹ Singer JE, Westphal M, Niswander K. Relationship of weight gain during pregnancy to birth weight and infant growth and development in the first year of life. *Obstetrics & Gynecology*. 1968 Mar;31(3):417-23.

²⁰² USDA Center for Nutrition Policy and Promotion. A look at the diet of pregnant women. *Nutrition Insights*, 17. 2000. Accessed February 23, 2009. www.cnpp.usda.gov/Publications/NutritionInsights/Insight17.pdf.

²⁰³ Wiles R. The views of women of above average weight about appropriate weight gain in pregnancy. *Midwifery*. 1998;14:254-260.

²⁰⁴ Fowles ER. Comparing pregnant women's nutritional knowledge to their actual dietary intake. *American Journal of Maternal Child Nursing*. 2002 May-Jun;27(3):171-177.

²⁰⁵ Lokey EA, Tran ZV, Wells CL, Myers BC, Tran AC. Effects of physical exercise on pregnancy outcomes: a meta-analytic review. *Medicine & Science in Sports & Exercise*. 1991 Nov;23(11):1234-9.

²⁰⁶ Clapp III JF. The effects of maternal exercise on early pregnancy outcome. *American Journal of Obstetrics & Gynecology*. 1989 Dec;161(6 Pt 1):1453-7.

²⁰⁷ Kulpa PJ, White BM, Visscher R. Aerobic exercise in pregnancy. *American Journal of Obstetrics & Gynecology*. 1987 Jun;156(6):1395-403.

²⁰⁸ Klebanoff MA, Shiono PH, Carey JC. The effect of physical activity during pregnancy on preterm delivery and birth weight. *American Journal of Obstetrics & Gynecology*. 1990 Nov;163(5 Pt 1):1450-6.

-
- ²⁰⁹ Hall DC, Kaufmann DA. Effects of aerobic and strength conditioning on pregnancy outcomes. *American Journal of Obstetrics & Gynecology*. 1987 Nov;157(5):1199-203.
- ²¹⁰ Hatch MC et al. Maternal exercise during pregnancy, physical fitness, and fetal growth. *American Journal of Epidemiology*. 1993 May 15;137(10):1105-14.
- ²¹¹ Clapp III JF, Simonian S, Lopez B, Appleby-Wineberg S, Harcar-Sevcik R. The one-year morphometric and neurodevelopmental outcome of the offspring of women who continued to exercise regularly throughout pregnancy. *American Journal of Obstetrics & Gynecology*. 1998 Mar;178(3):594-9.
- ²¹² Sternfeld B, Quesenberry CP, Jr., Eskenazi B, Newman LA. Exercise during pregnancy and pregnancy outcome. *Medicine & Science in Sports & Exercise*. 1995 May;27(5):634-40.
- ²¹³ Clapp III JF, Lopez B, Harcar-Sevcik R. Neonatal behavioral profile of the offspring of women who continued to exercise regularly throughout pregnancy. *American Journal of Obstetrics & Gynecology*. 1999 Jan;180(1 Pt 1):91-4.
- ²¹⁴ Kardel KR, Kase T. Training in pregnant women: effects on fetal development and birth. *American Journal of Obstetrics & Gynecology*. 1998 Feb;178(2):280-6.
- ²¹⁵ Davies GAL, Wolfe LA, Mottola MF, MacKinnon C. Society of Obstetricians and gynecologists of Canada SCPOC. Joint SOGC/CSEP clinical practice guideline: exercise in pregnancy and the postpartum period. *Canadian Journal of Applied Physiology*. 2003 Jun;28(3):330-41.
- ²¹⁶ Boscaglia N, Skouteris H, Wertheim, EH. Changes in body image satisfaction during pregnancy; a comparison of high exercising and low exercising women. *Australian and New Zealand Journal of Obstetrics and Gynaecology*. 2003;43:41-45.
- ²¹⁷ Symons Downs D, Hausenblas HA. Women's exercise beliefs and behaviors during their pregnancy and postpartum. *Journal of Midwifery and Women's Health*. 2004;49:138-144.
- ²¹⁸ Power ML, Cogswell ME, Schulkin J. Obesity prevention and treatment practices of US obstetrician-gynecologists. *Obstetrics & Gynecology*. 2006;108(4):961-968.
- ²¹⁹ Clarke PE, Gross H, Psychol C. Women's behaviour, beliefs and information sources about physical exercise in pregnancy. *Midwifery*. 2004; 20:133-141.
- ²²⁰ Devine CM, Bove CF, Olson CM. Continuity and change in women's weight orientations and lifestyle practices through pregnancy and the postpartum period: the influence of life course trajectories and transitional events. *Social Science and Medicine*. 2000;50:567-582.
- ²²¹ Wood Baker C, Carter AS, Cohen LR, Brownell KD. Eating attitudes and behaviors in pregnancy and postpartum global stability versus specific transitions. *Annals of Behavioral Medicine*. 1999;21(2):143-148.

- ²²² Polley BA, Wing RR, Sims CJ. Randomized controlled trial to prevent excessive weight gain in pregnant women. *International Journal of Obesity & Related Metabolic Disorders*. 2002 Nov;26(11):1494-502.
- ²²³ Olson CM, Strawderman MS, Reed RG. Efficacy of an intervention to prevent excessive gestational weight gain. *American Journal of Obstetrics & Gynecology*. 2004 Aug;191(2):530-6.
- ²²⁴ Claesson IM et al. Weight gain restriction for obese pregnant women: a case-control intervention study. *BJOG*. 2008 Jan;115(1):44-50.
- ²²⁵ Gray-Donald K, Robinson E, Collier A, David K, Renaud L, Rodrigues S. Intervening to reduce weight gain in pregnancy and gestational diabetes mellitus in Cree communities: an evaluation. *CMAJ*. 2000 Nov 14;163(10):1247-51.
- ²²⁶ Schmitt NM, Nicholson WK, Schmitt J. The association of pregnancy and the development of obesity – results of a systematic review and meta-analysis on the natural history of postpartum weight retention. *International Journal of Obesity*. 2007;31:1642-1651.
- ²²⁷ Keppel KG, Taffel SM. Pregnancy-related weight gain and retention: implications of the 1990 Institute of Medicine guidelines. *American Journal of Public Health*. 1993 Aug;83(8):1100-3.
- ²²⁸ Linne Y, Rossner S. Interrelationships between weight development and weight retention in subsequent pregnancies: the SPAWN study. *Acta Obstet Gynecol Scand*. 2003;82:318-25.
- ²²⁹ Gore SA, Brown DM, Smith West D. The role of postpartum weight retention in obesity among women: a review of the evidence. *Annals of Behavioral Medicine*. 2003;26(2):149-159.
- ²³⁰ Thorsdottir I, Birgisdottir BE. Different weight gain in women of normal weight before pregnancy: postpartum weight and birth weight. *Obstetrics & Gynecology*. 1998 Sep;92(3):377-83.
- ²³¹ Scholl TO, Hediger ML, Schall JI, Ances IG, Smith WK. Gestational weight gain, pregnancy outcome, and postpartum weight retention. *Obstetrics & Gynecology*. 1995 Sep;86(3):423-7.
- ²³² Kac G, Benicio MHDA, Velasquez-Melendez G, Valente JG, Struchiner CJ. Gestational weight gain and prepregnancy weight influence postpartum weight retention in a cohort of Brazilian women. *Journal of Nutrition*. 2004 Mar;134(3):661-6.
- ²³³ Olson CM, Strawderman MS, Hinton PS, Pearson TA. Gestational weight gain and postpartum behaviors associated with weight change from early pregnancy to 1 y postpartum. *International Journal of Obesity & Related Metabolic Disorders*. 2003 Jan;27(1):117-27.
- ²³⁴ Walker LO. Predictors of weight gain at 6 and 18 months after childbirth: a pilot study. *JOGNN - Journal of Obstetric, Gynecologic, & Neonatal Nursing*. 1996 Jan;25(1):39-48.

- ²³⁵ Ohlin A, Rossner S. Maternal body weight development after pregnancy. *International Journal of Obesity*. 1990 Feb;14(2):159-73.
- ²³⁶ Janney CA, Zhang D, Sowers M. Lactation and weight retention. *American Journal of Clinical Nutrition*. 1997 Nov;66(5):1116-24.
- ²³⁷ Walker LO et al. Weight and behavioral and psychosocial factors among ethnically diverse, low-income women after childbirth: I. Methods and context. *Women & Health*. 2004;40(2):1-17.
- ²³⁸ Fowles ER, Walker LO. Correlates of dietary quality and weight retention in postpartum women. *Journal of Community Health Nursing*. 2006;23(3):183-197.
- ²³⁹ Walker LO, Sterling BS, Timmerman GM. Retention of pregnancy-related weight in the early postpartum period: implications for women's health services. *JOGNN*. 2005;34:418-427.
- ²⁴⁰ Nuss H, Clarke K, Klohe-Lehman D, Freeland-Graves J. Influence of nutrition attitudes and motivators for eating on postpartum weight status in low-income new mothers. *Journal of the American Dietetic Association*. 2006;106:1774-1782.
- ²⁴¹ Pereira MA et al. Predictors of change in physical activity during and after pregnancy: project Viva. *American Journal of Preventive Medicine*. 2007;32(4):321-319.
- ²⁴² Gunderson EP et al. Association of fewer hours of sleep at 6 months postpartum with substantial weight retention at 1 year postpartum. *American Journal of Epidemiology*. 2008 Jan;167(2):178-87.
- ²⁴³ British Columbia Reproductive Care Program. Reproductive Mental Health Guideline 3: Identification and Assessment of Reproductive Mental Illness During the Preconception and Perinatal Periods. 2003. Accessed August 18, 2008. <http://www.bcphp.ca/sites/bcphp/files/Guidelines/Rmhg/Guideline3NewJan2003.pdf>
- ²⁴⁴ Carter AS, Wood Baker C, Brownell KD. Body Mass Index, eating attitudes, and symptoms of depression and anxiety in pregnancy and the postpartum period. *Psychosomatic Medicine*. 2000;62:264-270.
- ²⁴⁵ Fraser AB, Grimes DA. Effect of lactation on maternal body weight: a systematic review. *Obstet Gynecol Survey*. 2003;58(4):265-269.
- ²⁴⁶ Olson CM. Tracking of food choices across the transition to motherhood. *Journal of Nutrition Education and Behavior*. 2005;37:139-136.
- ²⁴⁷ Hill, JO. Obesity treatment: does one size fit all? *American Society for Clinical Nutrition*. 2005;81:1253-1254.
- ²⁴⁸ Harvey EL, Glennly A-M, Kirk SFL, Summberbell CD. Improving health professionals' management and the organization of care for overweight and obese people. *Cochrane Database of Systematic Reviews* 2001;2.

- ²⁴⁹ Gilden Tsai A, Wadden TA. Systematic review: an evaluation of major commercial weight loss programs in the United States. *Annals of Internal Medicine*. 2005;142:56-66.
- ²⁵⁰ Wilding JPH. Treatment strategies for obesity. *Obesity Reviews*. 2007;8(Suppl 1):137-144.
- ²⁵¹ Barclay L, Lie D, Martin BN. Behavioral intervention may help overweight, obese patients lost and maintain weight. *Journal of the American Medical Association*. 2008;299:1139-1148.
- ²⁵² Shaw K, O'Rourke P, Del Mar C, Kenardy, J. Psychological interventions for overweight or obesity. *Cochrane Database of Systematic Reviews*. 2005;2.
- ²⁵³ Wardle J. Eating behaviour and obesity. *Obesity Reviews*. 2007;8(Suppl 1):73-75.
- ²⁵⁴ Shaw K, O'Rourke P, Del Mar C. Exercise for overweight or obesity. *Cochrane Database of Systematic Reviews*. 2006;4.
- ²⁵⁵ Hill AJ. Obesity and eating disorders. *Obesity Reviews*. 2007;8(Suppl 1):151-155.
- ²⁵⁶ Brehm BJ et al. The role of energy expenditure in the differential weight loss in obese women on low fat and low carbohydrate diets. *The Journal of Clinical Endocrinology and Metabolism*. 2005;90:1475-1482.
- ²⁵⁷ Hare-Bruun H, Flint A, Heitmann BL. Glycemic index and glycemic load in relation to changes in body weight, body fat distribution, and body composition in adult Danes. *American Journal of Clinical Nutrition*. 2006;84(4):871-879.
- ²⁵⁸ Thomas DE, Elliot EJ, Baur L. Low glycaemic index or low glycaemic load diets for overweight and obesity. *Cochrane Database of Systemic Reviews*. 2007;3.
- ²⁵⁹ Bravata DM et al. Efficacy and safety of low-carbohydrate diets: a systematic review. *JAMA*. 2003;289(14):1837-1850.
- ²⁶⁰ Stern L et al. The effects of low-carbohydrate versus conventional weight loss diets in severely obese adults: one-year follow-up of a randomized trial. *Annals of Internal Medicine*. 2004;140(10):769-777.
- ²⁶¹ Amorim AR, Linne YM, Lourenco PMC. Diet or exercise, or both, for weight reduction in women after childbirth. *The Cochrane Collaboration*. 2007;4.
- ²⁶² Walker LO. Managing excessive weight gain during pregnancy and the postpartum period. *JOGNN*. 2007;36:490-500.
- ²⁶³ George GC, Hanss-Nuss H, Milani TJ, Freeland-Graves JH. Food choices of low-income women during pregnancy and postpartum. *Journal of the American Dietetic Association*. 2005;105:899-907.
- ²⁶⁴ Daley AJ, MacArthur C, Winter H. The role of exercise in treating postpartum depression: a review of the literature. *Journal of Midwifery and Women's Health*. 2007;52:56-62.

- ²⁶⁵ Clapp III J, Dickstein S. Endurance exercise and pregnancy outcome. *Medicine and Science in Sport and Exercise*. 1984;16:556-562.
- ²⁶⁶ Setse R, Grogan R, Cooper LA, Strobino D, Powe NR, Nicholson W. Weight loss programs for urban-based, postpartum African-American women: perceived barriers and preferred components. *Maternal Child Health Journal*. 2008;12:119-127.
- ²⁶⁷ Kinnunen TI, Pasanen M, Aittasalo M, Fogelholm M, Weiderpass E, Luoto R. Reducing postpartum weight retention – a pilot trial in primary health care. *Nutrition Journal*. 2007;6:21-29.
- ²⁶⁸ Koplan JP, Liverman CT, Kraak VI. Eds., Institute of Medicine. *Preventing Childhood Obesity: Health in the Balance*. Washington, DC: The National Academies Press;2005.
- ²⁶⁹ Birch LL, Fisher JO. Development of eating behaviors among children and adolescents. *Paediatrics*. 1998;101:539-549.
- ²⁷⁰ Spruijt-Metz D, Lindquist CH, Birch LL, Fisher JO, Goran MI. Relation between mothers' child-feeding practices and children's adiposity. *American Journal of Clinical Nutrition*. 2002;75:581-586.
- ²⁷¹ Wardle J, Carnell S, Cooke L. Parental control over feeding and children's fruit and vegetable intake: How are they related? *Journal of the American Dietetic Association*. 2005;105:227-232.
- ²⁷² Brown T, Kelly S, Summerbell C. Prevention of obesity: a review of interventions. *Obesity Reviews*. 2007;8(Suppl 1):127-130.
- ²⁷³ Robinson TN, Kiernan M, Matheson DM, Farish Haydel K. Is parental control of children's eating associated with childhood obesity? Results from a population-based sample of third graders. *Obesity Research*. 2001;9:306-312.
- ²⁷⁴ Lindsay AC, Sussner KM, Kim J, Gortmaker S. The role of parents in preventing childhood obesity. *Future of Children*. 2006;16(1):169-186.
- ²⁷⁵ Cullen KW et al. Availability, accessibility, and preferences for fruit, 100% fruit juice, and vegetables influence children's dietary behavior. *Health Education Behavior*. 2003;30(5):615-626.
- ²⁷⁶ Hood MY et al. Parental eating attitudes and the development of obesity in children: The Framingham Children's Study. *International Journal of Obesity and Related Metabolic Disorders*. 2000;24(10):1319-25.
- ²⁷⁷ Berry D et al. Family-based interventions for childhood obesity: a review. *Journal of Family Nursing*. 2004;10:429-449.
- ²⁷⁸ British Columbia Forum on Childhood Obesity. *Proceedings*. 2005.
- ²⁷⁹ Raine KD. *Overweight and Obesity in Canada: a population Health Perspective*. Ottawa: Canadian Institute for Health Information; 2004.

²⁸⁰ Caprio S. Treating child obesity and associated medical conditions. *Future of Children*. 2006;16(1):209-224.

²⁸¹ Gingras J. *Toxic Playground: Understanding the Environmental Influences that Increase the Risk of Childhood Weight Disturbance & Recommendations for Prevention*. South Fraser Health Region (now Fraser Health Authority); 2001.

²⁸² Registered Nurses' Association of Ontario. *Primary Prevention of Childhood Obesity; Nursing Best Practice Guideline*. 2005. Accessed February 23, 2009. www.rnao.org/bestpractices.

²⁸³ Ritchie L et al. *Prevention of Childhood Overweight – What Should Be Done? Position Paper*. Center for Weight and Health. U.C. Berkeley. 2001. Accessed February 23, 2009. http://cnr.berkeley.edu/cwh/PDFs/Prev_Child_Oweight_10-28-02.pdf

²⁸⁴ Satter EM. Internal regulation and the evolution of normal growth as the basis for prevention of obesity in children. *Journal of the American Dietetic Association*. 1996;96(9): 860-864.

²⁸⁵ Health Canada. *Nutrition for a Healthy Pregnancy: National Guidelines for the Childbearing Years*. Ottawa: Minister of Public Works and Government Services Canada; 1999.

²⁸⁶ Arendas K, Qiu Q, Gruslin A. Obesity in pregnancy: pre-conceptional to postpartum consequences. *J Obstet Gynaecol Can*. 2008 Jun;30(6):477-88.